

RESEARCH ARTICLE

10.1002/2016JD025190

Key Points:

- Two measures of growing season duration increased around western Lake Superior during 1984–2013
- Changes in lake temperature and ice cover regime strongly affect area T and P patterns and trends
- Changed regional T and P patterns and trends may be significant enough to modify forest phenology

Supporting Information:

- Supporting Information S1

Correspondence to:

M. Garcia,
matt.e.garcia@gmail.com

Citation:

Garcia, M., and P. A. Townsend (2016), Recent climatological trends and potential influences on forest phenology around western Lake Superior, USA, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2016JD025190.

Received 5 APR 2016

Accepted 5 NOV 2016

Accepted article online 9 NOV 2016

Recent climatological trends and potential influences on forest phenology around western Lake Superior, USA

Matthew Garcia¹ and Philip A. Townsend¹¹Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, Madison, Wisconsin, USA

Abstract We assess long-term climatological means, trends, and interannual variability around the western end of Lake Superior during 1984–2013 by using available weather station data. Our results focus on changes in basic and derived climate indicators from seasonal and annual temperature and precipitation, to the traditionally defined frost-free season, to a novel definition of the climatological growing season. We describe seasonal and year-to-year climate variability that influences forest phenology, using an alternative growing season metric that is based on the warm-season plateau in accumulated chilling days as an indicator of environmental triggers for vegetation growth and senescence. Our results indicate +0.56°C regional warming during our 30 year study period, with cooler springs (−1.26°C) and significant autumn warming (+1.54°C). The duration of the climatological growing season has increased +0.27 d/yr, extending primarily into autumn. Summer precipitation in our study area has declined by an average −0.34 cm/yr, potentially leading to moisture stress that impairs vegetation carbon uptake rates and can render the forest more vulnerable to disturbance. Many changes in temperature, precipitation, and climatological growing season are most prominent in locations where Lake Superior exerts a strong hydroclimatological influence, especially the Minnesota shoreline and in forest areas downwind (southeast) of the lake. Observed trends in lake temperature and ice phenology have also changed, coincident with a large-scale climatological regime shift around 1998. A number of factors are likely altering forest phenology and the role of the forest in the climate system of this ecologically important and highly varied forest-and-lake region.

1. Introduction

Changes in the climatology and related weather patterns of the Great Lakes region [Sousounis and Grover, 2002; Hayhoe et al., 2010; Huff and Thomas, 2014] are expected to affect both vegetation phenology [Jolly et al., 2005; Schwartz et al., 2006; Groffman et al., 2012; Gunderson et al., 2012; Jeong et al., 2013; Richardson et al., 2013] and forest disturbance regimes [Hufkens et al., 2012; Filewood and Thomas, 2014]. Forests exhibit both direct and long-term indirect feedback responses to changes in climate [Heide, 1993; Pope et al., 2013; Marchin et al., 2015], complicating our ability to predict forest phenology, estimate carbon sequestration, and represent clearly the numerous land-atmosphere interactions within the climate system [Peñuelas et al., 2009; Richardson et al., 2013]. Proximity to large water bodies such as Lake Superior can affect local and regional temperature and precipitation patterns [Changnon and Jones, 1972; Scott and Huff, 1996; Hinkel and Nelson, 2012]. The interactions of land-lake processes with climate change will be complex; spatiotemporal warming patterns and consequential changes in forest phenology may vary considerably across the Great Lakes region [Reyer et al., 2013]. In this paper we assess long-term climatological means, trends, and interannual variability in large forested areas around the western end of Lake Superior. We focus on describing changes in basic and derived climate indicators, from seasonal and annual temperature and precipitation to a novel definition of the climatological growing season.

Vegetation and forest phenology refer here to the seasonal timing of events in a tree's annual physiological cycle, including leaf budburst and growth, senescence, and (for deciduous species) abscission. Phenology also includes flowering and seed production, the generation of annual growth rings in stemwood, winter hardening and spring sap flush, and other less visible processes. From the tree's perspective, the timing of phenological events involves complex and dynamic trade-offs among numerous processes that are both driven and constrained by environmental conditions: photosynthesis balances carbon assimilation with moisture loss and nutrient transport; carbon must be allocated among leaf and stem growth, wood production [Delpierre et al., 2016], chemical defenses, and reproductive processes; and the tree must protect itself against temperature-related stresses such as freezing during spring and autumn transitions [Kim et al., 2014] and moisture-related stresses in hot and/or dry periods [Arend and Fromm, 2007; Anderegg et al., 2012]. The pace

of photosynthesis is acutely sensitive to environmental conditions, especially temperature [Ali *et al.*, 2015], and varies over time with the phenological phases of the leaf and tree. A tree must constantly adjust its allocation of resources (carbon, nutrients, moisture, and sunlight) according to changing biological strategies, subject to biological and abiotic limitations, in a competitive environment.

Forest phenology in our midlatitude temperate study area is driven largely by the annual temperature cycle, and phenological events can typically be ordered in “thermal time” by the accounting of cold season chilling days (CD) and warm season growing degree-days (GDD) [Baskerville and Emin, 1969; Cannell and Smith, 1983; Heide, 1993; Trudgill *et al.*, 2005]. Meteorological and climatological factors thus significantly influence vegetation seasonal phenology at the land surface [Jolly *et al.*, 2005; Ceccherini *et al.*, 2014; Hwang *et al.*, 2014; Koster *et al.*, 2014; Xie *et al.*, 2015] and explain a large fraction of observed year-to-year variability in temperate forest phenology [Fisher *et al.*, 2007; Marchin *et al.*, 2015], specifically as drivers of the annual growing season start, intensity, and duration.

Climate change in the Great Lakes region is expected to proceed at different rates for different seasons, with greater average warming expected in winter than in summer [Hayhoe *et al.*, 2007, 2010]. Precipitation regimes are expected to change, with a greater frequency of heavy precipitation events [Groisman *et al.*, 2012] and a diminishing proportion of winter precipitation falling as snow [Feng and Hu, 2007; Mishra and Cherkauer, 2011]. Warming winters may interfere with dormancy periods for species with leaf bud differentiation and development requirements [Rohde and Bhalerao, 2007; Morin *et al.*, 2009; Viherä-Aarnio *et al.*, 2014; Williams *et al.*, 2014], which can then affect the timing of spring budburst [Cannell and Smith, 1983; Murray *et al.*, 1989]. Young trees and undergrowth can be affected by changes in winter precipitation regimes and seasonal snow cover, especially where snowpack often insulates seedlings and soil from hard freezing [Drescher and Thomas, 2013].

A regional trend toward earlier spring green-up [Schwartz *et al.*, 2006; Morin *et al.*, 2009; Jeong *et al.*, 2011] can have substantial consequences for the ecosystem: overall seasonal carbon uptake and sequestration may increase [Saxe *et al.*, 2001; Millard *et al.*, 2007], but new growth is also exposed to an increased likelihood of spring frost events [Hänninen, 1991]. Some species may readily adapt to changing early-season freezing regimes [Saxe *et al.*, 2001] depending on the magnitude of interannual variability driving such changes. Winter warming along with greater variability in spring meteorological conditions may lead to more “false spring” and frost events that can damage leaves and severely hinder phenological processes through the remainder of the growing season [Rigby and Porporato, 2008; Augspurger, 2013; Peterson and Abatzoglou, 2014].

Uncertainty regarding climate change impacts on forests is even greater for autumn transitions, generally because the senescence process and its triggers remain poorly understood [Estiarte and Peñuelas, 2015; Gallinat *et al.*, 2015]. Autumn senescence and deciduous leaf abscission occur with photosynthetic downregulation [Hörtensteiner, 2006; Guo, 2013] and nutrient conservation as defenses against freezing injury [Killingbeck, 1996; Niinemets and Tamm, 2005]. A regional trend toward later leaf senescence [Jeong *et al.*, 2011] can lead to overall longer growing seasons [Jeong *et al.*, 2011; Gunderson *et al.*, 2012] and possibly to increased total primary productivity [Nemani *et al.*, 2003; Twine and Kucharik, 2009]. However, individuals and species for which phenological triggers adjust more slowly to environmental changes may remain susceptible to both frost and drought stress later in the growing season [Saxe *et al.*, 2001; Parida and Buermann, 2014]. Under drought conditions, a nutrient conservation process similar to winter preparation may drive leaf dormancy and senescence [Munné-Bosch and Alegre, 2004; Marchin *et al.*, 2010], and chronic moisture stress can compromise the long-term capacity for carbon assimilation in these forests [Noormets *et al.*, 2008; Brzostek *et al.*, 2014; Anderegg *et al.*, 2012, 2013, 2015].

Understanding climate change impacts on forest phenology requires both the comprehensive characterization of recent climatological variability, so that we can more accurately assess important trends, and an improved understanding of forest responses to that variability, which can differ across species and landscapes. In this paper we analyze spatiotemporal climate variability and trends during 1984–2013 in forests of the Upper Great Lakes, focusing primarily on proximity to Lake Superior and long-term climatological changes to the cold and warm seasons that may have had profound, observable effects on forest phenology in the region. We employ several threshold-based metrics in common use to characterize seasonal climatology: chilling days (CD) and freezing days (FD) from autumn through spring, and growing degree-days (GDD)




Figure 1. Study area examined in this work (dashed box) in regional context with collected GHCND surface weather stations, each indicated with its measurement type(s) and observation record duration.

from spring through autumn. We introduce an alternative definition of the climatological growing season based on the accumulation of CD through the year, similar but not equivalent to the more traditional frost-free growing season. Using these and several derivative metrics, we assess recent changes in regional climatology for the 1984–2013 period and examine spatiotemporal variability in growing season influences, identifying specific areas where changes and trends have differed markedly from others during our study period. This study presents an examination of the *climatological* growing season; our ongoing work seeks to associate that influence with the *observed* forest vegetation growing season across our study area. Follow-on work will apply our findings in this paper to an observational study of forest phenology over the same area and period using remote sensing methods.

2. Study Area and Background

Our study area covers $\sim 202,000 \text{ km}^2$ of subboreal evergreen and midlatitude mixed forest around the western end of Lake Superior in the North American Upper Great Lakes (Figure 1). This region hosts diverse forest and wildlife species, numerous protected and managed areas including state and national forests, widespread forest-related industry, and extensive tourism and recreational opportunities. A “tension zone” [Curtis and McIntosh, 1951] that traverses our study area is defined by a combination of geographic transitions, from warm and dry continental interior to cool and wet lake-influenced landscapes. This tension zone marks the approximate southern extent of the Laurentian glaciation with resulting gradients in soil types [Schaeztl *et al.*, 2005; Danz *et al.*, 2013] and encompasses a gradient in natural vegetation types [Wheeler *et al.*, 1992; Bockheim and Schliemann, 2014], from prairie and hardwood forests in the southwest (now mixed with agriculture) to subboreal evergreen and temperate mixed forests closer to Lake Superior. This transition is clearly visible in land cover maps of the region based on the U.S. Geological Survey National Land Cover Database products [Jin *et al.*, 2013; Homer *et al.*, 2015] (Figure S1 in the supporting information) and can be identified by using the U.S. Environmental Protection Agency [2011] ecoregion maps of our study area [Omernik *et al.*, 2000; Omernik, 2004] (Figure S2).

Our choice of study area reflects a particular challenge that arises from the complex geography of the region. Situated at the prairie-forest ecotone, the region borders the North American boreal forest but also has strong

agricultural influences, is subject to land-lake interactions and climatological influences of the Great Lakes. Forests in this region display a wide variety in observed phenology and disturbance events, with further variety expected as a consequence of climate change impacts on all of these elements. The forests of the Upper Great Lakes are increasingly vulnerable to disturbance factors and tree mortality due to seasonal moisture stress and fire risk under changing climatic conditions [Irland *et al.*, 2001]. This region is also sensitive to changing forest ecology and management practices, increasing recreational use and logging pressure, anthropogenic warming and climate change, and evolving conditions in the Great Lakes themselves including rapid recent warming of Lake Superior. While all of these factors may affect the forest and its role in the carbon cycle, in this paper we concentrate on the dominant climatological factors that may affect regional forest phenology and its interannual variability.

Upper Great Lakes climatology is influenced by the polar jet stream that frequently traverses the region and generally marks a continental-scale boundary between cold/dry polar air and warm/humid subtropical air along the primary midlatitude storm track. The jet stream over the region is directed by interactions between global circulations, synoptic dynamics, and climatological teleconnections [Rohli *et al.*, 1999; Grise *et al.*, 2013]. Both the upper level jet stream and surface land-lake interactions drive surface temperature gradients, frontal positions, and storm meteorology [Payer *et al.*, 2011]. Seasonality is a key factor in meteorological patterns at temperate latitudes, with synoptic variability dominating the spring and autumn transition seasons [Grover and Sousounis, 2002; Small and Islam, 2009; Small *et al.*, 2010] and the Great Lakes providing a strong regional influence on temperature and precipitation patterns throughout the year.

Given the position of the Great Lakes near the middle of the North American continent, jet stream and storm track patterns across the region are driven by numerous teleconnections including the Pacific-North America (PNA) pattern [Rodionov and Assel, 2001], the Arctic and North Atlantic Oscillations (AO and NAO, respectively) [Nie *et al.*, 2008; Luo and Cha, 2012], and often the combination of these with the Pacific Ocean El Niño–Southern Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO) [Bond and Harrison, 2000; Grise *et al.*, 2013]. Climatological teleconnections have been correlated most strongly with Great Lakes regional winter conditions [Rodionov and Assel, 2000, 2003; Wise *et al.*, 2015] and seasonal lake ice cover [Assel and Rodionov, 1998; Assel *et al.*, 2003; Bai *et al.*, 2011, 2012; Benson *et al.*, 2012; Bai and Wang, 2012; Wang *et al.*, 2012]. The warm phase of the long-period Atlantic Multidecadal Oscillation (AMO) has been associated with cold winters in eastern North America [Peings and Magnusdottir, 2014]. With a 4 to 7 year cycle, the ENSO cycle is often associated with warm/dry winters in the Upper Midwest U.S. during El Niño (warm) years and cool/wet winters during La Niña (cold) episodes [Trenberth *et al.*, 1998; McPhaden *et al.*, 2006].

Several studies have associated a climatological regime shift in the Upper Great Lakes region around 1998 associated with influential long- and short-period oscillating teleconnections: the AMO entered a primarily warm phase around 1995, and the PDO entered a persistently cold phase in 1998 coincident with an anomalously strong El Niño episode during the 1997–1998 winter [Bond *et al.*, 2003; Peterson and Schwing, 2003; Jo *et al.*, 2014]. Changes in Lake Superior ice cover, evaporation, water level, and water temperature regimes and trends have been identified around the same time [Van Cleave *et al.*, 2014; Watras *et al.*, 2014]. Great Lakes surface water temperatures have warmed faster than surrounding land surface temperatures [Austin and Colman, 2007, 2008; Van Cleave *et al.*, 2014] and Lake Superior ranks among the fastest-warming freshwater lakes in the world [O'Reilly *et al.*, 2015]. This warming has driven changes in the lake ice regime: peak areal coverage has decreased [Howk, 2009; Wang *et al.*, 2012], and observations indicate later ice onset in winter and breakup in spring [Assel, 2003; Howk, 2009; Assel *et al.*, 2013]. Reduced lake ice cover allows an overall increase in lake surface evaporation through the early winter [Brown and Duguay, 2010], resulting in greater winter lake-effect precipitation in downwind areas [Scott and Huff, 1996; Wright *et al.*, 2013].

3. Data and Methods

We developed climatological maps and statistics by using daily minimum and maximum temperature (T_{\min} and T_{\max}) and precipitation (P) observations from the merged Global Historical Climate Network–Daily (GHCND) data set [Durre *et al.*, 2010; Menne *et al.*, 2012] for 410 weather stations in the vicinity of western Lake Superior over the period from 1 January 1983 through 31 December 2013 (Figure 1). A number of these stations, including 83 stations in Canada and others outside of our analyzed portion of the U.S., are used here only for complete coverage of our study area. Our GHCND data set and the Python code used for our analyses

are available online at <https://megarcia.github.io/WxCD>. We relied on GHCND quality assurance flags and applied all valid station data for each date to generate gridded daily T_{\min} , T_{\max} , and P fields at 480 m spatial resolution by using radial basis functions [Akkala *et al.*, 2010]. This method in Python/SciPy is an exact multi-quadratic interpolator [Hardy, 1971, 1990; Franke *et al.*, 1994] with results that compare favorably with other spatial interpolators [Garcia *et al.*, 2008].

There is no explicit temporal interpolation of station values to cover periods of missing data at individual stations. As shown in Figure 1, some weather stations cover almost the entire analysis period, while many were active for only portions of that period. We sought to generate the best possible maps on a daily basis by using all available and valid station information for each day. All subsequent operations, including aggregation to seasonal averages or totals and statistical analyses for trends, are performed by using this series of gridded daily observations. We evaluated spatial interpolation error characteristics by a jackknife procedure (Figure S3) and found that our results are comparable to the Wisconsin-oriented climatological study by *Serbin and Kucharik* [2009]. Our spatial interpolation methods produced a daily T_{\min} bias (mean error) of $+0.022^{\circ}\text{C}$ and mean absolute errors (MAE) of 1.78°C , daily T_{\max} bias of $+0.038^{\circ}\text{C}$ and MAE of 1.60°C , and daily P bias of $+0.001$ cm and MAE of 0.152 cm. These bias values are less than the reporting precision of the meteorological stations in the GHCND data set (0.1°C and 0.025 cm for T and P , respectively), and our MAE values fall generally within expected interpolation accuracy [Garcia *et al.*, 2008].

We calculated the daily average temperature [Cannell and Smith, 1983] from the gridded fields as

$$T_{\text{avg}} = \frac{T_{\min} + T_{\max}}{2} \quad (1)$$

We then accumulated chilling days (CD), chilling degree-days (CDD), and growing degree-days (GDD) [de Reaumur, 1735; Baskerville and Emin, 1969; Thompson and Moncrieff, 1982; Lechowicz, 1984] on a daily basis as

$$\text{CD} = \sum_{i=0}^{\text{current date}} d_i \text{ with } d_i = \begin{cases} 1 & \text{if } T_{\text{avg},i} < T_{\text{base}} \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

$$\text{CDD} = \sum_{i=0}^{\text{current date}} dd_i \text{ with } dd_i = \max[0, (T_{\text{base}} - T_{\text{avg},i})] \quad (3)$$

$$\text{GDD} = \sum_{i=0}^{\text{current date}} dd_i \text{ with } dd_i = \max[0, (T_{\text{avg},i} - T_{\text{base}})] \quad (4)$$

We used $T_{\text{base}} = 5^{\circ}\text{C}$ following numerous empirical studies of tree physiology and spring phenology [Cannell and Smith, 1983; Murray *et al.*, 1989; Hunter and Lechowicz, 1992; Fisher *et al.*, 2007; Schenker *et al.*, 2014; Viherä-Aarnio *et al.*, 2014; Körner, 2015]. Accepted values of T_{base} vary strongly with species and setting, and those used for analysis of forest areas can differ greatly from values used in agricultural application [e. g., Skaggs and Irmak, 2012]. In general, there is an inverse relationship between the value of T_{base} and the accumulation of GDD required to attain certain phenological phases such as budburst or flowering [Trudgill *et al.*, 2005]. We also recognize that T_{base} and the biophysical efficiency of GDD and photoperiod use in early phenophases can vary widely among species and even within species across climatological settings. In the observational phase of our work, we may be able to evaluate such variety by using remote sensing indicators of phenological phases. For simplicity in this climatological analysis, and based on an example discussed by Trudgill *et al.* [2005], we have selected a single value of T_{base} in common use for temperate tree species to be applied across our study area. On this common basis, we can use calculated GDD accumulations with vegetation greenness observations that will be incorporated in follow-on work to differentiate among fast- and slow-developing, or more and less cold-adapted, phenological types.

We also calculated the accumulation of freezing days (FD, using $T_{\text{base}} = 0^{\circ}\text{C}$) through the cold season using both T_{avg} and T_{\min} similar to equation (2). Cold season variables (CD, CDD, and FD) are accumulated from 1 July through the following spring, and warm season variables (GDD) are accumulated from 1 January through the calendar year. We calculated seasonal temperature statistics as the 90 day mean and variance of T_{\min} , T_{\max} , and T_{avg} , total P over 90 day and longer periods up to a full year, and the accumulated days within a 90 day period with any ($P > 0$), moderate ($1 \text{ cm} < P \leq 2.5 \text{ cm}$), and heavy ($P > 2.5 \text{ cm}$) precipitation.

In total, we examined 66 climatological indicators (excluding teleconnection indices; see Appendix A) including several that have been used previously to assess changing climate extremes [Frich et al., 2002; Alexander et al., 2006].

We defined the vegetation growing season in two ways. The first is a traditional definition based on the time between last spring frost and first autumn frost dates ($T_{\min} < 0^{\circ}\text{C}$) [Kunkel et al., 2004; Skaggs and Irmak, 2012; Yu et al., 2014; McCabe et al., 2015]. Frost dates and the duration of the frost-free season are useful for assessing long-term seasonal changes as well as acute indicators of possible vegetation freeze damage, especially in the transition from cold to warm seasons [Augspurger, 2013]. To evaluate this danger we examined GDD accumulation at the date of the last spring frost. With more accumulated GDD near the beginning of the growing season, there is a greater chance that the opening of flowers and leaves on many species makes them vulnerable to a freezing event that can adversely affect tree productivity through the remainder of the growing season. In some cases, severe late frost events have followed a false spring period brought on by shifts in synoptic influences over several weeks, from cold to warm and then to cold again. Two such events within our study area and period, in 2007 and 2010, will be discussed below.

Alternatively, we also defined the growing season as a function of chilling day (CD) accumulation. We observed that CD accumulation reaches a warm-season plateau soon after GDD accumulation begins in the spring. The CD accumulation then departs from that plateau in the autumn just before GDD accumulation ceases. Between these dates, $T_{\text{avg}} \geq T_{\text{base}}$ and vegetation defenses against freezing are limited while net primary productivity is generally dedicated to growth and reproduction [Schenker et al., 2014; Vitasse et al., 2014; Körner, 2015; Pagter et al., 2015]. Our resulting plateau-based growing season is several days longer than the frost-free season for the same year. The differences between these season starting and ending dates are of great interest to us: in these periods, many forest species are vulnerable to environmental conditions that could affect vegetation carbon uptake over the entire growing season, as in the case of a late spring frost, or bring the growing season to an early close with an autumn frost event that triggers leaf senescence and the tree's winter preparations.

Using our alternative definition of the growing season, we evaluated two aggregate seasonal measures. We defined the "cold season intensity" as

$$\text{CSI} = \text{CDD}_{\text{plateau}} / \text{CD}_{\text{plateau}} \tag{5}$$

Cold season duration is represented well by CD, but that measure does not indicate the severity of the season: the winter could be particularly cold or relatively mild, but CD will continue to accumulate as long as $T_{\text{avg}} < T_{\text{base}}$. Conversely, accumulated CDD alone has little phenological meaning unless it is related to a calendar duration of some importance. Together, CD and CSI provide a composite indication of both the duration and severity of the cold season. Two winters with similar CD accumulations may be differentiated by their CSI values, providing a distinction between cold and mild seasons. Although trees may not recognize a difference between one subfreezing temperature and another, with regional warming we expect both CD and CSI (as calculated by our definitions) to diminish over time. Warmer winters can interfere with endodormancy ("winter chill") requirements for many forest species [Morin et al., 2009], leading to altered phenological cycles and reduced primary productivity in subsequent growing seasons.

Our aggregate measure for the warm season is the "growing season intensity" that we defined as

$$\text{GSI} = \frac{\text{GDD}_{\text{plateau end}} - \text{GDD}_{\text{plateau begin}}}{\text{DOY}_{\text{plateau end}} - \text{DOY}_{\text{plateau begin}}} \tag{6}$$

with DOY as the day of year and referring to the CD plateau as discussed above. In this case, both GDD and growing season duration remain relevant: phenological stages (e.g., budburst, leaf size thresholds, and maturity) are frequently associated with GDD accumulation [Cannell and Smith, 1983; Trudgill et al., 2005], and there is evidence that senescence timing depends on both inherent and external limits to leaf longevity, such as photoperiod [Kikuzawa et al., 2013]. Our GSI metric is separately useful as an indicator of temperature-related influences throughout the growing season, such as transpiration moisture demand and, if precipitation is inadequate through the season, the likelihood of vegetation moisture stress [Koster et al., 2014]. Tracking GSI through the growing season, especially using its rate of change on a short-term basis, enables




Figure 2. The 1984–2013 mean daily temperatures averaged over the U.S. portion of our study area, with observed seasonal indicators marked.

[Howk, 2009] and supplemented those with operation records from the nearby Madeline Island Ferry Line (MIFL) for 2011–2014 (Mary Ross, personal communication, 2015). Both of these sources indicate in their overlapping period that the area of Lake Superior near Bayfield did not completely freeze during the 2011–2012 winter season, and ice-on/ice-off dates otherwise differ between these sources by only 1–2 days; thus, we considered MIFL observations a reasonable proxy for NSIDC records that were not yet available for this work.

Finally, we evaluated the possible influences of global teleconnections on our regional climatology by using monthly indices reported by the NOAA National Centers for Environmental Prediction Climate Prediction Center and the NOAA Earth Systems Research Laboratory Physical Sciences Division. From these records we calculated 3 month averages for each season (December–January–February, March–April–May, June–July–August, and September–October–November) for each climate oscillation and pattern described in section 2 except ENSO. The ENSO signal is traditionally summarized over an extended winter (December–January–February–March) period, for which we calculated the average index value in each of the reported equatorial Pacific Ocean regions in order to identify any differences in their influence: Niño-3 (eastern), Niño-4 (central), and Niño-3.4 (overlapping parts of Niño-3 and Niño-4).

There are a number of caveats to be noted regarding the time series analyses that we perform here. Although the 30 year period that we have examined generally meets the customary duration criterion for climatological analyses, it is a relatively short period for trend analysis. Within our study period, we have assumed and analyzed trends by linear regression instead of higher-order functions that may fit the examined time series better. We ignore possible breaks and shifts in reporting long-term trends for our 30 year analysis. Although we do examine any changes in (linear) trends across the 1997–1998 winter season that can be extracted from these time series, there is a necessary trade-off where each of those trends covers a much shorter period and is less likely to represent a long-term climatological trend. We recognize that trends important to the physical system may not demonstrate statistical significance and that not all statistically significant trends indicate physical processes that can be justified in a conceptual model of the land-lake-atmosphere system examined here. We are particularly sensitive to the likelihood that derived climate indicators, such as plateau-based growing season duration, incorporate more basic variables that may demonstrate their own significant trends, even though a trend in the derived indicator may not itself end up being significant. Such cases can sometimes be attributed to opposing trends in the constituent indicators.

4. Results

Area-averaged 1984–2013 mean daily temperature observations (T_{min} , T_{max} , and T_{avg}) through the year are shown in Figure 2, which also provides a conceptual overview of key temperature-related climate variables that we analyze: CD, FD, and GDD accumulations; mean last spring and first autumn frost dates; and mean

subseasonal monitoring of vegetation status that can lead to reduced capacity for carbon uptake and growth [Teskey *et al.*, 1987], leaf wilting [Munné-Bosch and Alegre, 2004; Marchin *et al.*, 2010], and litter drying with evolving conditions conducive to forest fires [Yebra *et al.*, 2013] and other disturbance agents.

To examine the possible effects of Lake Superior seasonal ice phenology on nearby land areas, we included observations of ice onset and breakup at Bayfield, Wisconsin, from two sources. We obtained records for 1984–2012 from the NOAA National Snow and Ice Data Center (NSIDC)




Figure 3. The 1984–2013 mean CD and GDD accumulations averaged over the U.S. portion of our study area, with seasons marked.

concentrated on the Minnesota (northern) shore of Lake Superior, and the coldest temperatures in all seasons are found in the land areas northwest of the lake (Figures 5a, 5c, and 5e). Warmer temperatures in autumn and winter are generally concentrated on the Wisconsin/Michigan (southern) shore of Lake Superior, although some warm locations along the Minnesota shore are also apparent. The warmest temperatures in all seasons are found in the southwestern portion of our study area in primarily agricultural regions (Figure S1). The resulting spatial gradient in T_{avg} is strongest in a general north-south orientation across the far western end of Lake Superior and more generally across the tension zone that traverses our study area. Long-term positive trends in T_{min} (Figure 5b) are generally more widespread and stronger than those in T_{max} (Figure 5d), which actually demonstrates some areas of long-term cooling both northwest and southeast of Lake Superior over the study period.

From the area-average trends listed in Table 1 we calculated a 30 year net warming of $+0.56^{\circ}\text{C}$ in annual mean T_{avg} for our study area. Long-term cooling in spring (net $-1.26^{\circ}\text{C}/30$ years) has been offset by warming in all other seasons, especially autumn (net $+1.54^{\circ}\text{C}/30$ years), with the remainder of overall 30 year net warming divided almost evenly between the winter and summer seasons. Though none of these overall temperature trends demonstrate statistical significance at the $p < 0.05$ level using area-averaged values, significant trends can be found in more localized portions of our study area. In particular, long-term warming is evident in T_{min} (Figure 5b) and T_{avg} (Figure 5f) on the Minnesota shore of Lake Superior and following the approximate location of the tension zone across our study area.

CD plateau beginning and ending dates. The area-averaged mean annual accumulations of CD and GDD are shown in Figure 3 and are also marked with key dates related to the CD plateau. Area-averaged seasonal and annual precipitation (P) totals for 1984–2013 are shown in Figure 4. Though we refer here to seasonal T and P means and trends over our study period, maps of individual seasonal indicators are included in the supporting information that accompanies this paper. The area-averaged climatological mean, variance, trend, and extreme values for a number of both seasonal and annual climatological indicators are summarized in Table 1. Maps of study-period mean and trend values over 1984–2013 for several key climatological indicators are shown in Figures 5–10 with areas of trend significance at $p < 0.05$ marked.

4.1. Temperature and Precipitation Indicators and Lake Superior Influences

Seasonal and annual summaries of area-averaged T and P indicators are listed in the upper half of Table 1. Maps of annual average T values and interannual trends are shown in Figure 5. Cooler temperatures in spring and summer are concentrated




Figure 4. The 1984–2013 seasonal and annual precipitation averaged over the U.S. portion of our study area.

Maps of annual average *P* values and interannual trends are shown in Figure 6. Mean seasonal precipitation (Figure S7) is lowest in winter and highest in summer, with generally ambiguous area-wide spatial gradients in those seasons, while spatial gradients in mean seasonal *P* across our study area are generally north-to-south in spring and west-to-east in autumn, with comparable total *P* in those seasons. These patterns result

Table 1. 1984–2013 Climatological Indicator Statistics Averaged Over the U.S. Portion of Our Study Area^a

| Climatological Variable | Mean | SD | Trend (units/yr) | Minimum (Year) | Maximum (Year) |
|--|--------|-------|---------------------|---|-----------------------------|
| Temperature Indicators | | | | | |
| Winter average temperature (°C) | −9.6 | 2.3 | +0.04 | −13.7 (1994) | −4.8 (2012 [%]) |
| Spring average temperature (°C) | 9.2 | 1.4 | −0.04 | 6.7 (1996 [%]) | 11.6 (1987 [#]) |
| Summer average temperature (°C) | 17.7 | 0.9 | +0.03 | 15.1 (1992 [#]) | 19.4 (2002) |
| Autumn average temperature (°C) | 0.7 | 1.6 | +0.05 | −2.8 (1985 [%]) | 3.9 (2001 [%]) |
| Annual average temperature (°C) | 4.5 | 1.0 | +0.02 | 2.8 (1996 [%]) | 6.8 (1998 [#]) |
| Precipitation Indicators | | | | | |
| Winter total precipitation (cm) | 9.2 | 1.9 | +0.03 | 4.3 (1987 [#]) | 12.8 (1997) |
| Spring total precipitation (cm) | 23.4 | 5.1 | +0.13 | 14.1 (1988 [#]) | 36.4 (2001 [%]) |
| Summer total precipitation (cm) | 28.9 | 5.6 | −0.34 ^{**} | 18.4 (2012 [%]) | 39.6 (1999 [%]) |
| Autumn total precipitation (cm) | 17.0 | 4.9 | −0.02 | 9.7 (1989 [%]) | 26.6 (1996 [%]) |
| Annual total precipitation (cm) | 80.0 | 8.7 | −0.19 | 63.9 (2006 [%]) | 96.5 (1985) |
| Lake Superior Indicators | | | | | |
| Lake Superior ice-on date (DOY) | 15.9 | 18.7 | +1.15 ^{**} | −13 (1986/1990) | 69 (2002) |
| Lake Superior ice-off date (DOY) | 88.3 | 12.2 | +0.69 | 63 (2000 [%]) | 115 (1996 [%]) |
| Lake Superior ice duration (days) | 72.4 | 26.9 | −0.46 | 0 (1998 [#] /2012 [%]) | 109 (1996 [%]) |
| Cold Season Indicators | | | | | |
| Freezing days (using <i>T</i> _{min}) | 183.5 | 10.3 | −0.28 | 167.0 (2010 [#]) | 200.6 (1996 [%]) |
| Chilling days (using <i>T</i> _{avg}) | 176.9 | 10.5 | −0.14 | 158.6 (2012 [%]) | 196.3 (1996 [%]) |
| CSI (degrees) | 11.0 | 1.3 | −0.03 | 8.7 (2012 [%]) | 13.3 (1996 [%]) |
| Last spring freezing night (DOY) | 137.8 | 5.7 | −0.12 | 127.2 (2001 [%]) | 148.8 (1992 [#]) |
| First autumn freezing night (DOY) | 269.8 | 5.1 | +0.20 | 261.5 (1986) | 279.1 (2013) |
| Warm Season Indicators | | | | | |
| Last spring freezing night (GDD) | 155.6 | 47.0 | −2.12 [*] | 79.7 (1996 [%]) | 244.4 (1992 [#]) |
| Frost-free season (days) | 132.0 | 6.8 | +0.32 [*] | 116.9 (1992 [#]) | 144.5 (2005) |
| Beginning of CD plateau (DOY) | 127.4 | 6.8 | +0.16 | 112.7 (1985 [%]) | 139.7 (1992 [#]) |
| End of CD plateau (DOY) | 275.1 | 6.1 | +0.27 [*] | 262.9 (1991) | 286.6 (2013) |
| Plateau duration (days) | 147.7 | 7.3 | +0.11 | 134.4 (1991) | 165.5 (1998 [#]) |
| Plateau GDD (degree-days) | 1659.6 | 134.5 | +1.88 | 1316.4 (1992 [#]) | 1896.8 (1988 [#]) |
| GSI (degrees) | 11.2 | 0.8 | +0.00 | 9.5 (1992 [#]) | 13.0 (1991) |

^aTrend significance is marked as * at *p* < 0.05 and ** at *p* < 0.01. Extreme years affected by El Niño (ENSO+) and La Niña (ENSO−) events are marked with # and %, respectively.




Figure 5. The 1984–2013 mean annual T_{min} , T_{max} , and T_{avg} , with trends. Areas of trend significance at $p < 0.05$ are stippled.

in a southeastward gradient in mean annual P that is most prominent across Lake Superior (Figure 6a). We note here the lake-effect influence on spatial distributions of P in winter, with the areas of heaviest winter precipitation on the southern shore and immediately downwind (southeast) of Lake Superior. Winter precipitation is increasing in some areas, especially in the immediate vicinity of Lake Superior, but summer precipitation exhibits an area-average trend of -0.34 cm/yr ($p < 0.01$), with areas of stronger trends along almost the entire lakeshore and extending in a number of directions from Lake Superior, especially along




Figure 6. The 1984–2013 mean annual P , with trends. Areas of trend significance at $p < 0.05$ are stippled.

the tension zone in Wisconsin. Area-average reductions in summer precipitation are consistent with an overall trend in summer precipitation days ($P > 0$) of -0.25 d/yr ($p < 0.001$). The resulting trends in total annual precipitation (Figure 6b) are relatively neutral across much of our study area, with isolated locations of significant negative trends in areas where the summer trends dominate, especially south and southeast of Lake Superior.

A number of our results are consistent with the thermal influence of Lake Superior on regional climatological indicators. We found higher winter and lower summer temperatures along the northern lakeshore than in the forest region farther to the northwest, with a longer frost-free season along the lakeshore by more than 4 days. Growing season transitions, as indicated by the beginning and end of the CD plateau, are also delayed 2–3 days along the lakeshore in comparison with the forest areas to the northwest. The accumulation of




Figure 7. The 1984–2013 mean-derived cold season indicators, with trends. Areas of trend significance at $p < 0.05$ are stippled.




Figure 8. The 1984–2013 mean frost-based growing season start, end, and duration, with trends. Areas of trend significance at $p < 0.05$ are stippled.

area-wide winter precipitation days is positively correlated with the date of Lake Superior ice-on conditions ($p < 0.05$) and negatively correlated with ice cover duration ($p < 0.05$). The number of spring moderate precipitation days ($1 \text{ cm} < P \leq 2.5 \text{ cm}$) in areas downwind (south and southeast) of Lake Superior is negatively correlated with ice duration ($p < 0.05$), a result consistent with ice cover inhibiting lake evaporation and thus lake-effect precipitation in those areas.




Figure 9. The 1984–2013 mean plateau-based growing season start, end, and duration, with trends. Areas of trend significance at $p < 0.05$ are stippled.

Our results suggest that the seasonal ice cover on western Lake Superior plays an important role in temperature and precipitation patterns on adjacent land areas. Our analyses indicated a trend in the date of Lake Superior ice-on conditions of $+1.15 \text{ d/yr}$ ($p < 0.01$), although we recognize that a linear trend cannot adequately account for the large interannual variability in lake ice phenology. Notably, during autumn and winter the area of greatest precipitation in our study area is concentrated southeast of Lake Superior where prevailing northwesterly winds carry lake-evaporated moisture onshore. Conversely, spring and summer P appears




Figure 10. The 1984–2013 mean plateau-based growing season indicators, with trends. Areas of trend significance at $p < 0.05$ are stippled.

relatively suppressed in the same area. Cold-season ice cover from midwinter through early spring inhibits lake evaporation and lake effect precipitation; during the summer the lake is generally cooler than the surrounding land areas, inducing a stable atmospheric boundary layer and inhibiting the development of storms that could then move onshore [Changnon and Jones, 1972]. Even the small trends in lake ice phenology that we found are enough to allow statistically significant increases in winter precipitation (Figure S7) southeast of Lake Superior. However, spring precipitation seems also to increase in those same areas, which we cannot attribute to the slightly later lake ice breakup over time. In that case, our analyses may require a finer temporal division of seasonal precipitation in areas downwind of the lake so that portions of the spring season before and after the observed ice-off date in each year are treated separately. Such a division could help demonstrate the transition from winter (cold) to spring (warm) weather patterns and phenological processes in the area of lake influence.

4.2. Cold and Warm Season Indicators and Interseasonal Variability

As expected, the greatest accumulation of CD in our study area occurs in the forest areas northwest of Lake Superior (Figure 7a). The 30 year trend in CD accumulation (Figure 7b) is generally mixed across the study area, with only small areas of statistically significant reduction in CD accumulation that are generally coincident with the locations of winter T_{\min} and annual T_{avg} warming. There is slightly greater spatial variability evident in the detail of 30 year mean CSI (Figure 7c) which is lowest in the immediate vicinity of Lake Superior and in downwind regions. This result, and decreasing CSI over time in many areas on the lakeshore (Figure 7 d), is consistent with the warming surface temperatures and changing ice phenology of Lake Superior.

The duration of the frost-free season (Figure 8f), our traditional measure of the climatological growing season, shows an area-average trend of $+0.32 \text{ d/yr}$ ($p < 0.05$) over our 30 year study period (Table 1) that is consistent




Figure 11. The 1984–2013 intraseasonal and interseasonal temperature and precipitation correlations. Only correlations at $p < 0.05$ are shown, with * at $p < 0.01$ and ** at $p < 0.001$.

with previous studies [Easterling, 2002; Kunkel et al., 2004; Yu et al., 2014; McCabe et al., 2015]. There is much spatial detail in maps of trends in the last spring and first autumn frost dates (Figures 8b and 8d, respectively), with some areas experiencing statistically significant changes in both indicators consistent with lengthening of the frost-free season. Most of these areas occur in the forested areas northwest of Lake Superior and along the tension zone south and southeast of the lake.

We found quite different patterns for the CD plateau (Figure 9), our alternative measure of the climatological growing season. The beginning of the CD plateau changed little over time in the area average but shows a large area downwind of Lake Superior with significant trends toward a later start to the climatological growing season (Figure 9b). The end of the CD plateau showed an area-average trend of +0.27 d/yr ($p < 0.05$) over our 30 year study period with large areas of significant trends (Figure 9d) that are consistent with significant autumn warming. These include areas on the Minnesota (northern) shore of Lake Superior and across agricultural areas in the southwest portion of our study area, but notably not in the region immediately southeast of the lake. Trends in the duration of the CD plateau are strongly mixed across our study area (Figure 9f), with an overall longer plateau evident in many areas where the end of the plateau now extends further into autumn, but a shorter plateau in areas directly upwind and downwind of Lake Superior. Our maps of plateau GDD accumulation, GSI, and their trends (Figure 10) are consistent with slightly warmer climatological growing seasons over time, especially for isolated areas in immediate proximity to the lake.

We found throughout the year that seasonal mean average temperature is inversely correlated with the number of seasonal precipitation days ($P > 0$) but not with accumulated seasonal P , demonstrating that cloudy seasons are cooler seasons overall. We also found several significant ($p < 0.05$) interseasonal correlations among T and P indicators (Figure 11), especially between winter and spring conditions. Winter T and P indicators are strongly tied to T and the number of precipitation days in the following spring. However, by these methods we found no statistically significant interseasonal correlations that might be used to extend T and P predictability beyond spring and through the remainder of the growing season.

4.3. Teleconnection Influences and a Climatological Regime Shift

Two large-scale teleconnections examined here, the AMO and PDO, demonstrated regime transitions around the middle of our study period in the course of their long-period oscillations. NOAA observations indicated that the AMO index increased during 1984–2013 and around 1995 shifted to a warm (AMO+) phase that is generally associated with drier conditions in the Upper Midwest U.S. [Enfield et al., 2001]. The PDO index generally decreased through our 30 year study period, with observations of distinct PDO+ periods (indicating warmer northeastern Pacific coastal waters) in the mid-1980s and mid-1990s followed by PDO anomalies

(cool northeastern Pacific coastal waters) beginning around 1998. Our results indicate that the AMO (PDO) index is positively (negatively) correlated with study area T , winter P , and the date of Lake Superior ice onset. The AMO (PDO) index is negatively (positively) correlated with summer P and lake ice duration. Local correlations with Pacific teleconnections (PDO, ENSO, and PNA) appear strongest in winter and spring, with Pacific indices overall positively correlated with T and negatively correlated with P in the Upper Great Lakes region. A long-term increase in autumn PNA index values ($p < 0.05$), and positive values of the PNA index, in general, has been associated with relatively dry conditions over continental North America [Leathers *et al.*, 1991]. The often-related AO and NAO indices showed no significant correlation with study area surface climatology except in autumn, when the AO index is negatively correlated with area-averaged P .

A distinct shift in climatological regimes across several global and regional indicators was thus observed around 1998. Our analyses indicated changes to, and even reversals of, statistically significant ($p < 0.05$) temporal trends for several climatological indicators between the 1984–1998 and 1998–2013 periods. Specifically, paired maps of trends in spring and autumn T_{avg} and annual P are shown in Figure 12. Many locations with strong spring cooling during 1984–1998 (Figure 12a) shifted to near-neutral trends in the latter half of our study period (Figure 12b), which can be attributed primarily to a shift in the trend of spring T_{max} with very little change in spring T_{min} trends across much of our study area. Locations with weak autumn warming during the earlier period (Figure 12c) shifted to strong cooling trends in the 1998–2013 period (Figure 12d), which we attribute to sharp reversals in both autumn T_{max} and autumn T_{min} trends across our study area. Despite this reversal to statistically significant cooling in many locations, our strongest 30 year area-averaged seasonal warming still appears in autumn (Figure S6 and Table 1). Annual total P shifted from increasing trends in isolated locations during the 1984–1998 period (Figure 12e) to widespread decreasing trends across much of our study area in the 1998–2013 period (Figure 12f). We attribute this shift primarily to an area-average negative trend in summer P ($p < 0.01$; Figure S7 and Table 1) that is also strongest in the latter half of our study period and to a lesser extent diagnosed shifts to decreasing seasonal precipitation trends that we also found for winter and spring.

5. Discussion

Using available weather station data, we have examined the mean seasonal and annual climatology, temporal trends, teleconnection correlations, and the potential influences of Lake Superior on surrounding land areas during 1984–2013. Our study area is characterized by an extensive and ecologically important forest-and-lake landscape at the Upper Midwest U.S. prairie-forest ecotone, where we anticipate observable sensitivity to recent and ongoing climate change. Our results indicate regional warming of $+0.56^{\circ}\text{C}$ through our 30 year study period supported by mixed seasonal changes, with spring area-average cooling offset primarily by autumn warming. Long-term trends include warmer winters, wetter winter and spring seasons, a diminishing duration of Lake Superior ice cover, and a strong decrease in summer P since the 1998 PDO regime shift and El Niño event.

Using both traditional and new definitions of the growing season, we have identified an overall extension of the climatological growing season into autumn. Our new definition, based on a warm season plateau in CD accumulation, provides additional information for diagnosing early-season freezing risks and later periods of potential vegetation moisture stress, both of which can have a strong influence on seasonal and interannual variabilities in vegetation phenology. However, some risks such as late spring frost and false spring events may remain location-specific and are not characterized well with area-averaged metrics.

Several prominent global and hemispherical climatic teleconnections correlate in varying degrees with seasonal and interannual variabilities in regional hydroclimatology and with lake ice phenology, all of the factors affecting seasonal T and P patterns in our study area. Changes in Lake Superior ice cover regimes have contributed to greater winter and spring lake-effect P in portions of our study area south and southeast of the lake. Lake Superior clearly affects T and P patterns in nearby land areas through both proximity and an interseasonal lag due to thermal inertia, with effects that are likely strong enough to alter forest phenology in those areas.

5.1. Climatological and Growing Season Trends

The overall area-average temperature increase that we estimated by linear regression is consistent with prior estimates of warming in this region over similar periods [Li *et al.*, 2010; Groisman *et al.*, 2012]. Our results show a consistent continuation of trends reported for 1951–1980 climatic changes in the Great Lakes region




Figure 12. The 1984–1998 and 1998–2013 trends in selected climatological indicators.

reported by *Scott and Huff* [1996]: increased minimum temperatures in all seasons, decreased spring and summer maximum temperatures, slight decreases in summer rainfall, and large winter precipitation increases for lake-effect areas primarily southeast of Lake Superior and as winter ice cover diminished. *Choi et al.* [2014] analyzed *Serbin and Kucharik's* [2009] climatological data set for Wisconsin and showed statistically significant decreases in the frequency of heavy precipitation events during 1950–2006, especially southeast of Lake Superior in the same lake-effect areas where we also found slight decreases for heavy precipitation days ($P > 2.5$ cm) in summer and autumn. However, our analyses indicated slight increases in winter and spring

heavy precipitation days in those areas. These differences with *Choi et al.* [2014] might be attributed to differing study periods but are likely related to technical differences (e.g., study area boundaries, station data selection, interpolation methods, and grid resolution) in the examined precipitation fields that can demonstrate large spatial variability.

Our estimated $+0.56^{\circ}\text{C}$ area-average temperature change, along with the accelerated warming of Lake Superior surface waters ($+2.5^{\circ}\text{C}$) [*Austin and Colman*, 2007, 2008; *Van Cleave et al.*, 2014], clearly indicates regional warming over the 30 year study period. Though the beginning of the CD plateau in spring has changed little, we have noted the extension of the CD plateau later into autumn consistent with long-term temperature increases concentrated primarily in that season. Overall, these results point to a longer climatological growing season due more to warm-season extension later into autumn than earlier into spring, consistent with findings by *Jeong et al.* [2011]. Annual total precipitation demonstrated a slight negative area-average trend over the study period and a particularly strong negative trend since 1998 (Figure 12) driven by a sharp decrease in summer precipitation (Figure S7). Our analyses of temperature trends support the small negative trends that we found for cold season CD and FD and a consequent increase in the duration of the frost-free season. Along with observed influences of large-scale teleconnections (AMO+, PDO–, and increasing PNA), these changes suggest an overall drying trend [*Parida and Buermann*, 2014] within our study period for the region.

Area-wide climatological warming and drying during 1984–2013 means that forests in our study area may have experienced increasing moisture stress in that period, especially in summer. It is important here to distinguish between changes affecting moisture availability, based primarily on P , and those that drive vegetation moisture demand, based primarily on T . Vegetation moisture stress occurs with the combination of these, when moisture availability is insufficient to meet moisture demand. Conditions promoting moisture stress can inhibit transpiration and reduce growth [*Teskey et al.*, 1987], promote leaf wilt and early senescence [*Munné-Bosch and Alegre*, 2004; *Marchin et al.*, 2010], enhance tree mortality [*Anderegg et al.*, 2012, 2013], and reduce overall forest carbon uptake [*Brzostek et al.*, 2014; *Koster et al.*, 2014].

For tree species that adapt quickly to changing climate conditions and phenological cues, a longer climatological growing season may drive changes in growing season net primary productivity [*Nemani et al.*, 2003; *Twine and Kucharik*, 2009]. A warmer spring is typically associated with an earlier start to the vegetation growing season, provided adjustment to the timing and speed of leaf growth (subject to freezing risks) to these warmer conditions. Higher autumn temperatures may support longer vegetation growing seasons for those forest species that can adjust their leaf longevity and senescence triggers to warmer autumn conditions. However, we are still learning to identify and understand the many cues for leaf senescence that influence leaf phenology, including photoperiod and temperature [*Kikuzawa et al.*, 2013; *Ali et al.*, 2015] and biochemical limits on leaf longevity [*Keenan and Richardson*, 2015; *Seki et al.*, 2015].

5.2. Seasonal Transitions

Climate during the shoulder seasons can be critical to vegetation phenology. A lack of statistically significant interseasonal correlations between summer and the shoulder seasons (Figure 11) leads to uncertainty regarding aspects of forest phenology such as the progression of green-up, seasonal peak greenness in early summer, and deciduous autumn senescence with follow-on effects for the related winter feeding patterns of overwintering herbivores. Summer and autumn conditions strongly influence seed production in many forest species, with consequences for wildlife feeding and reproduction patterns [*Yang et al.*, 2010]. These growing season conditions affect biochemical processes that control leaf bud set and hardening prior to late autumn freezing, with consequences for optimum productivity in the next growing season [*Vitasse et al.*, 2014; *Estiarte and Peñuelas*, 2015]. Seasonal climatological conditions can thus affect forest phenology and overall primary productivity in the same season and well into the forest life cycle [*Noormets et al.*, 2008; *Anderegg et al.*, 2012, 2013, 2015; *Brzostek et al.*, 2014].

Regarding late frost events and false spring occurrences, we found an area-average trend of -2.1 degree-days/yr ($p < 0.05$) in the accumulation of GDD before the last spring frost. This trend, along with a slight trend toward earlier last spring freezing nights (Table 1), suggests a slowly diminishing risk of vegetation-damaging spring frost events over our study period. This result is not necessarily consistent with the frequency-based examination of frost-based spring vegetation damage [*Augsburger*, 2013] that suggests an increasing

frequency of occurrences in the Midwest US. We take particular note of false spring and late frost events that were observed in the U.S. Upper Midwest in 2007 [Augsburger, 2009; Gu et al., 2008] and 2010 [Fereday et al., 2012; Hufkens et al., 2012; Filewood and Thomas, 2014; Ning and Bradley, 2014]. For these years the area-average accumulations of GDD before the last spring frost were 147 and 156 degree-days, respectively, both very close to the 30 year average for that metric and with earlier-than-normal last spring frost dates in both years. By contrast, a spring heatwave in 2012 [Ellwood et al., 2013; Peterson et al., 2013] led to an area-average accumulation of 200 GDD by the time of the last spring frost, which also occurred earlier than the 30 year normal date, yet no particular frost-related damage was reported across the region. Hypothetically, we expect higher-than-normal GDD accumulations (suggesting greater likelihood of early leaf growth) before a possibly later-than-normal last spring frost date for these events. However, false springs in ENSO-neutral 2007 and El Niño-influenced 2010, but not in La Niña-influenced 2012, ran contrary to that expectation. Under that hypothesis, we further suspect that false spring events occurred for portions of our study area in 1986 and 1992, as indicated in Table 1. We anticipate that our continuing observational study of regional forest phenology will provide us with greater insight into these occurrences, given the potential importance of spring late frost events to seasonal phenological patterns and their impact on tree growth.

The seasonal and interseasonal climatological relationships that we have found may promote some interesting phenological patterns. Within a season, the inverse relationship between average temperature and precipitation days (as a proxy for cloudy days) corresponds to days with greater water availability (from precipitation) also having lower evaporation and transpiration moisture demand. A cycle may be established in which forest soil and canopy moisture are allowed to build up during periods with cloudy/wet days and then deplete during periods with clear/dry days. Lower leaf primary productivity in cloudy periods may then be offset by enhanced productivity during clear periods when supported by greater soil moisture availability. Increased soil moisture is also conducive to seed germination and seedling growth, especially for shade-tolerant species, and can thus have potential impacts on mixed forest understory structure and species distributions [Nowacki and Abrams, 2008].

Conversely, dry soils and forest litter can lead to inhibited soil respiration, slower litter decay, and potential interference with seed germination and seedling growth, also altering forest structure and composition over time [Gustafson and Sturtevant, 2013; Peters et al., 2015]. Reduced moisture availability can eventually lead to a shift of the surface energy balance away from latent heating (evaporation and transpiration) in the growing season to greater sensible heating at the surface and in the forest canopy, a positive feedback cycle that can enhance local warming and exacerbate forest canopy moisture stress [Anderegg et al., 2012]. A warmer spring and earlier start to the growing season may therefore compensate for a cooler summer in some years in terms of total growing season primary productivity, but the reverse is not necessarily true: a warm summer may not compensate for a cool spring and a late start to the growing season, but instead exacerbate temperature and moisture stresses that reduce forest productivity throughout the growing season. Numerous influences generate complex interactions around the Upper Great Lakes, and the lakes themselves contribute to spatiotemporal variability in the system through their internal mixing regimes, providing some “memory” of conditions across seasons (via thermal inertia) and potentially over several years [Bennett, 1978; Gerten and Adrian, 2001; Piccolroaz et al., 2015].

5.3. Teleconnection Influences

We are concerned not only with the mean climatology but also its year-to-year variability and extremes, as these also influence forest phenology and growth [Bouriaud et al., 2005; Voelker et al., 2012]. We may generalize for the Upper Midwest U.S. that El Niño (ENSO+) events foster warm/dry years and La Niña (ENSO−) events support cool/wet years: Table 1 lists extremes in many of our climatological indicators that can be associated with ENSO events within our study period. El Niño conditions that persisted through 1987 and 1988 produced the driest winter and warmest growing season in our study period in conjunction with regional drought conditions [Trenberth et al., 1988; Weaver et al., 2009]. On the other hand, Lake Superior did not fully freeze during both the El Niño winter of 1997–1998 [Changnon, 1999] and the La Niña winter of 2011–2012 [Peterson et al., 2013; Dole et al., 2014], the latter event leading into one of the most spatially and temporally variable growing seasons during our study period [Hoerling et al., 2014]. Teleconnection variability may have influenced the wettest year of our study period in 1985 (ENSO−, AMO−, and PDO+) and the driest year in 2006 (ENSO−, AMO+, and PDO−), both La Niña years. It remains to be seen whether our observed

teleconnection influences are part of a persistent long-term trend or might change, such as with eventual regime shifts in the long-period AMO and PDO.

Understanding interannual climatological variability, combined with interactions of the land-atmosphere system, in our study area is clearly more complex than reliance on any single element such as teleconnections [Wise *et al.*, 2015] or another external indicator will allow. As another example, Weaver *et al.* [2009] analyzed the 1993 Upper Midwest summer flood event, which was supported by both AO– and NAO– conditions, and found that moisture from the distant Gulf of Mexico contributed significantly to large rainfall totals in the Great Lakes and Upper Mississippi River regions that year. An event or season can reach climatological extremes by the interaction and reinforcement of influences: the NAO phase, which can drive synoptic organization over the Great Lakes but may not necessarily generate a significant event on its own, interacted with the alignment of AO-driven meridional flows to drive the northward transport of Gulf moisture along the entire length of the Mississippi River valley.

5.4. Lake Effects

Climatological characteristics are distinguished across the western end of Lake Superior and reflect the influence of the lake on T (northwestern shore) and P (to the southeast), both of which can be linked to the thermal inertia of Lake Superior surface waters during seasonal transition periods. Warm lake surface waters in autumn and winter enhance the land-lake temperature contrast, contributing also to strong P gradients across the lake from lower accumulations in the northwest to higher totals to the southeast. Delayed ice formation in warm winters allows a longer period of surface evaporation, feeding lake-effect P maxima to the southeast through both autumn and winter. Conversely, Lake Superior ice cover that extends well into spring in some years may contribute to diminished spring P as well as delayed last spring frost dates for that area. Because of the thermal and moisture effects of the lake, we anticipate that vegetation phenological transitions in the spring season for areas south and east of Lake Superior are typically delayed several days, possibly weeks, compared with those transitions west and north of the lake.

The influence of Lake Superior and its own changes over time on the climatology of our study area is substantial, including both recent lake warming [Austin and Colman, 2007, 2008; Van Cleave *et al.*, 2014; O'Reilly *et al.*, 2015] and changing ice phenology [Assel, 2003; Howk, 2009; Wang *et al.*, 2012; Assel *et al.*, 2013]. Observational studies regarding the effects of the Great Lakes on their surrounding land areas [e.g., Li *et al.*, 2010] remain essential to our growing understanding of land-atmosphere processes in the surrounding forest areas. Given their prevalence on the study area landscape (Figure S1), the roles of smaller lakes in regional climatology and forest phenology are also of interest to our work [Johnson and Stefan, 2006; Plank and Shuman, 2009; Mishra and Cherkauer, 2011; Mishra *et al.*, 2011a, 2011b]. Overall, while we remain interested in phenological events in the autumn season and the potential effects of Lake Superior thermal inertia on delayed timing of those events in the vicinity of the lake, the spring transition and its complexity because of that proximity are vital for understanding phenological transitions at the start of the vegetation growing season and will be an interesting point of focus in future work.

5.5. Potential Applications

Our analyses indicate that climatological influences are strongly heterogeneous across our study region but may be relatively homogeneous, or at least similar, over smaller distances on the order of 100 km. An analysis of climate trends in this area requires detailed examination of patterns at the subregional scale. We cannot assume that climatology and its influences on forest areas southeast of Lake Superior are similar to those in areas northwest of the lake or even that climatology is consistent throughout the nearby ecoclimatic tension zone. Along the northern shore of Lake Superior in Minnesota, an analysis of climatological influences on forest phenology and health must include localized details. Different regimes of land-atmosphere interaction among these regions may require differing approaches to the identification of dominant influences on forest phenology, the causes of forest disturbances, and the processes involved in postdisturbance forest recovery. Recent and continued climatic changes may promote altered trajectories of forest health (e.g., due to moisture stress and other disturbance factors) and forest species composition (favoring more temperate and drought-resistant species) [Duveneck *et al.*, 2014a, 2014b], complicating options for regional forest management toward long-term goals [Rittenhouse and Rissman, 2015]. Trophic interactions between forest vegetation, insects, and wildlife become more uncertain with altered timing of phenological events [Foster *et al.*,

2013; Roberts *et al.*, 2015]. Forest managers with responsibility for planning winter and dry-season harvest territories, or for undertaking intervention activities to mitigate forest disturbances due to insect pests and fire, can benefit from any potential improvements in seasonal predictability.

We seek a better understanding of recent climate and forest phenological variability to support the development and validation of land surface models that incorporate both permanent and transient land cover change as well as seasonal vegetation processes [White *et al.*, 1997]. There is an increasing need for improved accuracy in representation of land surface states and processes as regional and global climate models progress from relatively coarse ($\Delta x = 10\text{--}100$ km [Prein *et al.*, 2015]) to finer representative scales. As we develop a better understanding of this coupled system at a range of spatial and temporal scales, we will enable the capability to model feedback between forest phenology and climate change at local scales [Peñuelas *et al.*, 2009; Richardson *et al.*, 2013; Brzostek *et al.*, 2014], essential for accurate assessments of forest carbon states and their spatiotemporal variability [Desai *et al.*, 2008; Jeong *et al.*, 2013; Schwartz *et al.*, 2013; Lu *et al.*, 2015].

In regions such as the Great Lakes, we must exercise caution: most land-atmosphere models do not yet incorporate even large lakes in the physical system, producing often significant errors in the simulation of temperature and moisture fluxes [Bryan *et al.*, 2015]. A number of researchers have pursued better parameterizations of lake processes for use in energy- and water-balance models [Plank and Shuman, 2009; Mishra *et al.*, 2010] and for proper representation of lakes in regional climate models [Gula and Peltier, 2012; Bennington *et al.*, 2014; Mallard *et al.*, 2014]. It is important that large and small lakes, their surface temperatures and winter ice phenology, and their influences on surrounding areas are properly included in analyses of climate change effects on temperature and precipitation patterns and trends, so that regional modeling efforts [Hayhoe *et al.*, 2010; Gula and Peltier, 2012; Mallard *et al.*, 2014; Harding and Snyder, 2015; Notaro *et al.*, 2015] can provide more accurate insight and guidance to scientists and decision-makers.

6. Conclusions

A wide variety of environmental factors contribute to seasonal and year-to-year variabilities in forest phenology around the Upper Great Lakes. In this work we have identified several trends showing recent and likely ongoing changes to the climatological growing season and the availability of moisture to regional forests. Our results also show large spatial variability in trends such as autumn and winter warming, diminishing summer precipitation, and the extension of the climatological growing season into autumn. Dry summers can lead to moisture stress that impairs carbon uptake rates and may render the forest more vulnerable to disturbance factors.

Our new definition of the climatological growing season, based on a warm-season plateau in accumulated chilling days, provides information that can be combined with observations of the traditional frost-free growing season to show the speed, and potential dangers to vegetation, of the transition seasons. The timing and rates of spring green-up and autumn senescence can have large variability and have significant impacts on the seasonal carbon assimilation capacity of affected forest areas. The extension of warm periods into autumn can alter the environmental cues for leaf senescence and the preparation of trees for winter conditions, potentially impacting phenology in the following growing season as well. It is important to note that while meteorology changes with the season, and climate factors change from year to year, trees in the forest maintain memory of these conditions through the impacts on phenology and growth in subsequent seasons and thus over a lifetime.

Adding to the complexity of the study area, Lake Superior exerts an identifiable influence on surrounding and especially downwind land areas. The warming lake, with its changing winter ice phenology, contributes to winter warming in the immediate vicinity of the lakeshore. One of the greatest potential influences of Lake Superior on regional phenology is the contribution of longer ice-free periods in autumn and winter to lake-effect precipitation in large forest areas south and southeast of the lake. The resulting extension of cold periods later into spring may delay spring green-up in those areas at the time of year when seasonal leaf expansion and wood production are vital to the relative competitive advantages of different species in mixed forests, affecting both carbon sequestration and resource optimization to sustain optimum growth rates through the remainder of the warm season. Forest management in the context of climate change must therefore consider the range of factors affecting the critical green-up and senescence periods bounding the growing season.

Finally, our work provides valuable support for an observational effort to understand long-term changes in vegetation phenology including seasonal environmental cues and the impacts of climatological variability on forest phenology, disturbances, and postdisturbance forest recovery. All of these processes inform our developing understanding of forest responses to recent and ongoing climate change. Feedback from surface conditions (including lakes, land cover, land use, snow cover, vegetation phenology, and forest disturbances) to the climate system [e.g., Sobolowski *et al.*, 2010; Richardson *et al.*, 2013; Rydzik and Desai, 2014] are integral to the coevolution of the land-atmosphere system and thus important to any examination of regional phenoclimatology.

Appendix A

Seasonal and annual climatological indicators examined in this work. Traditional seasons are defined as 90 day periods ending on the equinox and solstice dates.

Temperature indicators (from spatially interpolated GHCND station data records)

1. Winter mean and temporal standard deviation of T_{min} , T_{max} , and T_{avg} (°C)
2. Spring mean and temporal standard deviation of T_{min} , T_{max} , and T_{avg} (°C)
3. Summer mean and temporal standard deviation of T_{min} , T_{max} , and T_{avg} (°C)
4. Autumn mean and temporal standard deviation of T_{min} , T_{max} , and T_{avg} (°C)
5. Annual (365 day) mean and temporal standard deviation of T_{min} , T_{max} , and T_{avg} (°C)

Precipitation indicators (from spatially interpolated GHCND station data records)

1. Winter total precipitation (cm) and precipitation days ($P > 0$)
2. Winter moderate ($1 \text{ cm} < P \leq 2.5 \text{ cm}$) and heavy ($P > 2.5 \text{ cm}$) precipitation days
3. Spring total precipitation (cm) and precipitation days ($P > 0$)
4. Spring moderate ($1 \text{ cm} < P \leq 2.5 \text{ cm}$) and heavy ($P > 2.5 \text{ cm}$) precipitation days
5. Summer total precipitation (cm) and precipitation days ($P > 0$)
6. Summer moderate ($1 \text{ cm} < P \leq 2.5 \text{ cm}$) and heavy ($P > 2.5 \text{ cm}$) precipitation days
7. Autumn total precipitation (cm) and precipitation days ($P > 0$)
8. Autumn moderate ($1 \text{ cm} < P \leq 2.5 \text{ cm}$) and heavy ($P > 2.5 \text{ cm}$) precipitation days
9. Annual (365 day) total precipitation (cm) (evaluated on 31 December)

Lake Superior (from NSIDC/MIFL records)

1. Ice-on and ice-off dates (day of year)
2. Ice duration (days)

Cold season indicators (derived from daily gridded data set)

1. CD and CDD at beginning of plateau (accumulated from 1 July using T_{avg})
2. FD at beginning of plateau (accumulated from 1 July using T_{min})
3. Cold season intensity (CSI, calculated by using plateau CDD and CD, in °C)
4. Date (day of year) of last spring freeze/frost (using T_{min})
5. Date (day of year) of first autumn freeze/frost (using T_{min})

Warm season indicators (derived from daily gridded data set)

1. Duration of frost-free season (days)
2. Date (day of year) at beginning of CD plateau (using T_{avg})
3. Date (day of year) at end of CD plateau (using T_{avg})
4. Duration of CD plateau (days)
5. GDD at date of beginning of CD plateau (accumulated from 1 January by using T_{avg})
6. GDD at date of last spring freeze/frost (accumulated from 1 January by using T_{avg})
7. GDD accumulated during spring and summer (using T_{avg})
8. GDD at date of end of CD plateau (accumulated from 1 January by using T_{avg})
9. GDD accumulated during CD plateau (using T_{avg})
10. Growing season intensity (GSI, calculated from plateau GDD and duration, in °C)

Acknowledgments

We acknowledge Ankur Desai's comments regarding an early draft of this work. Comments and suggestions by two anonymous reviewers significantly improved our manuscript. Support for this work was provided by the USDA Forest Service–Northern Research Station, USDA NIFA McIntire–Stennis funding to the University of Wisconsin–Madison (project WIS01621), UW–Madison College of Agricultural and Life Sciences, UW–Madison Department of Forest and Wildlife Ecology, NASA Biodiversity and Ecological Forecasting Project NNX14AC36G, and the first author's 2015–2016 Wisconsin Space Grant Consortium Graduate Research Fellowship. This work employed the computing resources and assistance of the UW–Madison Center for High Throughput Computing (CHTC) in the Department of Computer Sciences. The CHTC is supported by UW–Madison, the Advanced Computing Initiative, the Wisconsin Alumni Research Foundation, the Wisconsin Institutes for Discovery, and the National Science Foundation and is an active member of the Open Science Grid supported by the National Science Foundation and the U.S. Department of Energy's Office of Science. Our GHCND station data set and Python analysis code are described in the supporting information and available online at <https://megarcia.github.io/WxCD>.

References

- Akkala, A., V. Devabhaktuni, and A. Kumar (2010), Interpolation techniques and associated software for environmental data, *Environ. Prog. Sust. Energy*, *29*, 134–141, doi:10.1002/ep.10455.
- Alexander, L. V., et al. (2006), Global observed changes in daily climate extremes of temperature and precipitation, *J. Geophys. Res.*, *111*, D05109, doi:10.1029/2005JD006290.
- Ali, A. A., et al. (2015), Global-scale environmental control of plant photosynthetic capacity, *Ecol. Appl.*, *25*, 2349–2365, doi:10.1890/14-2111.1.
- Anderegg, W. R. L., J. M. Kane, and L. D. L. Anderegg (2012), Consequences of widespread tree mortality triggered by drought and temperature stress, *Nat. Clim. Change*, *3*, 30–36, doi:10.1038/nclimate1635.
- Anderegg, W. R. L., L. Plavcová, L. D. L. Anderegg, U. G. Hacke, J. A. Berry, and C. B. Field (2013), Drought's legacy: Multiyear hydraulic deterioration underlies widespread aspen forest die-off and portends increased future risk, *Global Change Biol.*, *19*, 1188–1196, doi:10.1111/gcb.12100.
- Anderegg, W. R. L., et al. (2015), Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models, *Science*, *349*, 528–532, doi:10.1126/science.aab1833.
- Arend, M., and J. Fromm (2007), Seasonal change in the drought response of wood cell development in poplar, *Tree Physiol.*, *27*, 985–992, doi:10.1093/treephys/27.7.985.
- Assel, R., and S. Rodionov (1998), Atmospheric teleconnections for annual maximum ice cover on the Laurentian Great Lakes, *Int. J. Climatol.*, *18*, 425–442, doi:10.1002/(SICI)1097-0088(19980330)18:4<425::AID-JOC258>3.0.CO;2-Q.
- Assel, R., K. Cronk, and D. Norton (2003), Recent trends in Laurentian Great Lakes ice cover, *Clim. Change*, *57*, 185–204, doi:10.1023/A:1022140604052.
- Assel, R. A. (2003), Great Lakes ice cover, first ice, last ice, and ice duration: Winters 1973–2002. NOAA Technical Memorandum GLERL-125, Great Lakes Environmental Research Laboratory, Ann Arbor, Michigan.
- Assel, R. A., J. Wang, A. H. Clites, and X. Bai (2013), Analysis of Great Lakes ice cover climatology: Winters 2006–2011. NOAA Technical Memorandum GLERL-157, Great Lakes Environmental Research Laboratory, Ann Arbor, Michigan.
- Augsburger, C. K. (2009), Spring 2007 warmth and frost: Phenology, damage and refoliation in a temperate deciduous forest, *Funct. Ecol.*, *23*, 1031–1039, doi:10.1111/j.1365-2435.2009.01587.x.
- Augsburger, C. K. (2013), Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing, *Ecology*, *94*, 41–50, doi:10.1890/12-0200.1.
- Austin, J., and S. Colman (2008), A century of temperature variability in Lake Superior, *Limnol. Oceanogr.*, *53*, 2724–2730, doi:10.4319/lo.2008.53.6.2724.
- Austin, J. A., and S. M. Colman (2007), Lake Superior summer water temperatures are increasing more rapidly than regional air temperatures: A positive ice-albedo feedback, *Geophys. Res. Lett.*, *34*, L06604, doi:10.1029/2006GL029021.
- Bai, X., and J. Wang (2012), Atmospheric teleconnection patterns associated with severe and mild ice cover on the Great Lakes, 1963–2011, *Water Qual. Res. J. Can.*, *47*, 421–435, doi:10.2166/wqrjc.2012.009.
- Bai, X., J. Wang, Q. Liu, D. Wang, and Y. Liu (2011), Severe ice conditions in the Bohai Sea, China, and mild ice conditions in the Great Lakes during the 2009/10 winter: Links to El Niño and a strong negative Arctic oscillation, *J. Appl. Meteorol. Climatol.*, *50*, 1922–1935, doi:10.1175/2011JAMC2675.1.
- Bai, X., J. Wang, C. Sellinger, A. Clites, and R. Assel (2012), Interannual variability of Great Lakes ice cover and its relationship to NAO and ENSO, *J. Geophys. Res.*, *117*, C03002, doi:10.1029/2010JC006932.
- Baskerville, G. L., and P. Emin (1969), Rapid estimation of heat accumulation from maximum and minimum temperatures, *Ecology*, *50*, 514–517, doi:10.2307/1933912.
- Bennett, E. B. (1978), Characteristics of the thermal regime of Lake Superior, *J. Great Lakes Res.*, *4*, 310–319, doi:10.1016/S0380-1330(78)72200-8.
- Bennington, V., M. Notaro, and K. D. Holman (2014), Improving climate sensitivity of deep lakes within a regional climate model and its impact on simulated climate, *J. Clim.*, *27*, 2886–2911, doi:10.1175/JCLI-D-13-00110.1.
- Benson, B. J., J. J. Magnuson, O. P. Jensen, V. M. Card, G. Hodgkins, J. Korhonen, D. M. Livingstone, K. M. Stewart, G. A. Weyhenmeyer, and N. G. Granin (2012), Extreme events, trends, and variability in Northern Hemisphere lake-ice phenology (1855–2005), *Clim. Change*, *112*, 299–323, doi:10.1007/s10584-011-0212-8.
- Bockheim, J. G., and S. A. Schliemann (2014), Soil richness and endemism across an environmental transition zone in Wisconsin, USA, *Catena*, *113*, 86–94, doi:10.1016/j.catena.2013.09.011.
- Bond, N. A., and D. E. Harrison (2000), The Pacific decadal oscillation, air-sea interaction and central north Pacific winter atmospheric regimes, *Geophys. Res. Lett.*, *27*, 731–734, doi:10.1029/1999GL010847.
- Bond, N. A., J. E. Overland, M. Spillane, and P. Stabeno (2003), Recent shifts in the state of the North Pacific, *Geophys. Res. Lett.*, *30*(23), 2183, doi:10.1029/2003GL018597.
- Bouriaud, O., J.-M. Leban, D. Bert, and C. Deleuze (2005), Intra-annual variations in climate influence growth and wood density of Norway spruce, *Tree Physiol.*, *25*, 651–660, doi:10.1093/treephys/25.6.651.
- Brown, L. C., and C. R. Duguay (2010), The response and role of ice cover in lake-climate interactions, *Prog. Phys. Geogr.*, *34*, 671–704, doi:10.1177/0309133310375653.
- Bryan, A. M., A. L. Steiner, and D. J. Posselt (2015), Regional modeling of surface-atmosphere interactions and their impact on Great Lakes hydroclimate, *J. Geophys. Res. Atmos.*, *120*, 1044–1064, doi:10.1002/2014JD022316.
- Brzostek, E. R., D. Dragoni, H. P. Schmid, A. F. Rahman, D. Sims, C. A. Wayson, D. Johnson, and R. P. Phillips (2014), Chronic water stress reduces tree growth and the carbon sink of deciduous hardwood forests, *Global Change Biol.*, *20*, 2531–2539, doi:10.1111/gcb.12528.
- Cannell, M. G. R., and R. I. Smith (1983), Thermal time, chill days and prediction of budburst in *Picea sitchensis*, *J. Appl. Ecol.*, *20*, 951–963, doi:10.2307/2403139.
- Ceccherini, G., N. Gobron, and M. Migliavacca (2014), On the response of European vegetation phenology to hydroclimatic anomalies, *Remote Sens.*, *6*, 3143–3169, doi:10.3390/rs6043143.
- Changnon, S. A. (1999), Impacts of 1997–98 El Niño generated weather in the United States, *Bull. Am. Meteorol. Soc.*, *80*, 1819–1827, doi:10.1175/1520-0477(1999)080<1819:IOENOG>2.0.CO;2.
- Changnon, S. A., Jr., and D. M. A. Jones (1972), Review of the influences of the Great Lakes on weather, *Water Resour. Res.*, *8*, 360–371, doi:10.1029/WR008i002p0360.
- Choi, W., R. Tareghian, J. Choi, and C.-S. Hwang (2014), Geographically heterogeneous temporal trends of extreme precipitation in Wisconsin, USA during 1950–2006, *Int. J. Climatol.*, *34*, 2841–2852, doi:10.1002/joc.3878.

- Curtis, J. T., and R. P. McIntosh (1951), An upland forest continuum in the prairie-forest border region of Wisconsin, *Ecology*, *32*, 476–496, doi:10.2307/1931725.
- Danz, N. P., L. E. Frelich, P. B. Reich, and G. J. Niemi (2013), Do vegetation boundaries display smooth or abrupt spatial transitions along environmental gradients? Evidence from the prairie–forest biome boundary of historic Minnesota, USA, *J. Veg. Sci.*, *24*, 1129–1140, doi:10.1111/jvs.12028.
- Delpierre, N., D. Berveiller, E. Grandia, and E. Dufréne (2016), Wood phenology, not carbon input, controls the interannual variability of wood growth in a temperate oak forest, *New Phytol.*, *210*, 459–470, doi:10.1111/nph.13771.
- de Reaumur, R. A. F. (1735), Observations du thermometre, faites à Paris pendant l'année 1735, comparées avec celles qui ont été faites sous la Ligne, à l'Isle de France, à Alger, et en quelques-unes de nos Isles de l'Amerique. *Mem. Acad. Royale des Sci.*, Paris, 545–576.
- Desai, A. R., et al. (2008), Influence of vegetation and seasonal forcing on carbon dioxide fluxes across the Upper Midwest, USA: Implications for regional scaling, *Agric. For. Meteorol.*, *148*, 288–308, doi:10.1016/j.agrformet.2007.08.001.
- Dole, R., et al. (2014), The making of an extreme event: Putting the pieces together, *Bull. Am. Meteorol. Soc.*, *95*, 427–440, doi:10.1175/BAMS-D-12-00069.1.
- Drescher, M., and S. C. Thomas (2013), Snow cover manipulations alter survival of early life stages of cold-temperate tree species, *Oikos*, *122*, 541–554, doi:10.1111/j.1600-0706.2012.20642.x.
- Durre, I., M. J. Menne, B. E. Gleason, T. G. Houston, and R. S. Vose (2010), Comprehensive automated quality assurance of daily surface observations, *J. Appl. Meteorol. Climatol.*, *49*, 1615–1633, doi:10.1175/2010JAMC2375.1.
- Duveneck, M. J., R. M. Scheller, M. A. White, S. D. Handler, and C. Ravenscroft (2014a), Climate change effects on northern Great Lake (USA) forests: A case for preserving diversity, *Ecosphere*, *5*, 23, doi:10.1890/ES13-00370.1.
- Duveneck, M. J., R. M. Scheller, and M. A. White (2014b), Effects of alternative forest management on biomass and species diversity in the face of climate change in the northern Great Lakes region (USA), *Can. J. Forest Res.*, *44*, 700–710, doi:10.1139/cjfr-2013-0391.
- Easterling, D. R. (2002), Recent changes in frost days and the frost-free season in the United States, *Bull. Am. Meteorol. Soc.*, *83*, 1327–1332, doi:10.1175/1520-0477(2002)083<1327:RCIFDA>2.3.CO;2.
- Ellwood, E. R., S. A. Temple, R. B. Primack, N. L. Bradley, and C. C. Davis (2013), Record-breaking early flowering in the eastern United States, *PLoS One*, *8*, e53788, doi:10.1371/journal.pone.0053788.
- Enfield, D. B., A. M. Mestas-Núñez, and P. J. Trimble (2001), The Atlantic multidecadal oscillation and its relation to rainfall and river flows in the continental U.S., *Geophys. Res. Lett.*, *28*, 2077–2080, doi:10.1029/2000GL012745.
- Estiarte, M., and J. Peñuelas (2015), Alteration of the phenology of leaf senescence and fall in winter deciduous species by climate change: Effects on nutrient proficiency, *Global Change Biol.*, *21*, 1005–1017, doi:10.1111/gcb.12804.
- Feng, S., and Q. Hu (2007), Changes in winter snowfall/precipitation ratio in the contiguous United States, *J. Geophys. Res.*, *112*, D15109, doi:10.1029/2007JD008397.
- Fereday, D. R., A. Maidens, A. Arribas, A. A. Scaife, and J. R. Knight (2012), Seasonal forecasts of northern hemisphere winter 2009/10, *Environ. Res. Lett.*, *7*, 034031, doi:10.1088/1748-9326/7/3/034031.
- Filewood, B., and S. C. Thomas (2014), Impacts of a spring heat wave on canopy processes in a northern hardwood forest, *Global Change Biol.*, *20*, 360–371, doi:10.1111/gcb.12354.
- Fisher, J. I., A. D. Richardson, and J. F. Mustard (2007), Phenology model from surface meteorology does not capture satellite-based greenupt estimations, *Global Change Biol.*, *13*, 707–721, doi:10.1111/j.1365-2486.2006.01311.x.
- Foster, J. R., P. A. Townsend, and D. J. Mladenoff (2013), Mapping asynchrony between gypsy moth egg-hatch and forest leaf-out: Putting the phenological window hypothesis in a spatial context, *Forest Ecol. Manage.*, *287*, 67–76, doi:10.1016/j.foreco.2012.09.006.
- Franke, R., H. Hagen, and G. M. Nielson (1994), Least squares surface approximation to scattered data using multiquadratic functions, *Adv. Comput. Math.*, *2*, 81–99, doi:10.1007/BF02519037.
- Frich, P., L. V. Alexander, P. Della-Marta, B. Gleason, M. Haylock, A. M. G. Klein Tank, and T. Peterson (2002), Observed coherent changes in climatic extremes during the second half of the twentieth century, *Clim. Res.*, *19*, 193–212, doi:10.3354/cr019193.
- Gallinat, A. S., R. B. Primack, and D. L. Wagner (2015), Autumn, the neglected season in climate change research, *Trends Ecol. Evol.*, *30*, 169–176, doi:10.1016/j.tree.2015.01.004.
- Garcia, M., C. D. Peters-Lidard, and D. C. Goodrich (2008), Spatial interpolation of precipitation in a dense gauge network for monsoon storm events in the southwestern United States, *Water Resour. Res.*, *44*, W05S13, doi:10.1029/2006WR005788.
- Gerten, D., and R. Adrian (2001), Differences in the persistency of the North Atlantic Oscillation signal among lakes, *Limnol. Oceanogr.*, *46*, 448–455, doi:10.4319/lo.2001.46.2.0448.
- Grise, K. M., S.-W. Son, and J. R. Gyakum (2013), Intraseasonal and interannual variability in North American storm tracks and its relationship to equatorial Pacific variability, *Mon. Weather Rev.*, *141*, 3610–3625, doi:10.1175/MWR-D-12-00322.1.
- Groffman, P. M., et al. (2012), Long-term integrated studies show complex and surprising effects of climate change in the northern hardwood forest, *BioScience*, *62*, 1056–1066, doi:10.1525/bio.2012.62.12.7.
- Groisman, P. Y., R. W. Knight, and T. R. Karl (2012), Changes in intense precipitation over the central United States, *J. Hydrometeorol.*, *13*, 47–66, doi:10.1175/JHM-D-11-039.1.
- Grover, E. K., and P. J. Sousounis (2002), The influence of large-scale flow on fall precipitation systems in the Great Lakes basin, *J. Clim.*, *15*, 1943–1956, doi:10.1175/1520-0442(2002)015<1943:TIOLSF>2.0.CO;2.
- Gu, L., P. J. Hanson, W. Mac Post, D. P. Kaiser, B. Yang, R. Nemani, S. G. Pallardy, and T. Meyers (2008), The 2007 eastern US spring freeze: Increased cold damage in a warming world?, *BioScience*, *58*, 253–262, doi:10.1641/B580311.
- Gula, J., and W. R. Peltier (2012), Dynamical downscaling over the Great Lakes basin of North America using the WRF regional climate model: The impact of the Great Lakes system on regional greenhouse warming, *J. Clim.*, *25*, 7723–7742, doi:10.1175/JCLI-D-11-00388.1.
- Gunderson, C. A., N. T. Edwards, A. V. Walker, K. H. O'Hara, C. M. Campion, and P. J. Hanson (2012), Forest phenology and a warmer climate—Growing season extension in relation to climatic provenance, *Global Change Biol.*, *18*, 2008–2025, doi:10.1111/j.1365-2486.2011.02632.x.
- Guo, Y. (2013), Towards systems biological understanding of leaf senescence, *Plant Mol. Biol.*, *82*, 519–528, doi:10.1007/s11103-012-9974-2.
- Gustafson, E. J., and B. R. Sturtevant (2013), Modeling forest mortality caused by drought stress: Implications for climate change, *Ecosystems*, *16*, 60–74, doi:10.1007/s10021-012-9596-1.
- Hänninen, H. (1991), Does climatic warming increase the risk of frost damage in northern trees?, *Plant, Cell Environ.*, *14*, 449–454, doi:10.1111/j.1365-3040.1991.tb01514.x.
- Harding, K. J., and P. K. Snyder (2015), The relationship between the Pacific–North American teleconnection pattern, the Great Plains low-level jet, and North Central U.S. heavy rainfall events, *J. Clim.*, *28*, 6729–6742, doi:10.1175/JCLI-D-14-00657.1.
- Hardy, R. L. (1971), Multiquadric equations of topography and other irregular surfaces, *J. Geophys. Res.*, *76*, 1905–1915, doi:10.1029/JB076i008p01905.

- Hardy, R. L. (1990), Theory and applications of the multiquadric–biharmonic method, *Comput. Math. Appl.*, *19*, 163–208, doi:10.1016/0898-1221(90)90272-L.
- Hayhoe, K., et al. (2007), Past and future changes in climate and hydrological indicators in the U.S. Northeast, *Clim. Dyn.*, *28*, 381–407, doi:10.1007/s00382-006-0187-8.
- Hayhoe, K., J. VanDorn, T. Croley II, N. Schlegal, and D. Wuebbles (2010), Regional climate change projections for Chicago and the US Great Lakes, *J. Great Lakes Res.*, *36*(Suppl. 2), 7–21, doi:10.1016/j.jglr.2010.03.012.
- Heide, O. M. (1993), Dormancy release in beech buds (*Fagus sylvatica*) requires both chilling and long days, *Physiol. Plant.*, *89*, 187–191, doi:10.1111/j.1399-3054.1993.tb01804.x.
- Hinkel, K. M., and F. E. Nelson (2012), Spatial and temporal aspects of the lake effect on the southern shore of Lake Superior, *Theor. Appl. Climatol.*, *109*, 415–428, doi:10.1007/s00704-012-0585-2.
- Hoerling, M., J. Eischeid, A. Kumar, R. Leung, A. Mariotti, K. Mo, S. Schubert, and R. Seager (2014), Causes and predictability of the 2012 Great Plains drought, *Bull. Am. Meteorol. Soc.*, *95*, 269–282, doi:10.1175/BAMS-D-13-00055.1.
- Homer, C. G., J. A. Dewitz, L. Yang, S. Jin, P. Danielson, G. Xian, J. Coulston, N. D. Herold, J. D. Wickham, and K. Megown (2015), Completion of the 2011 National Land Cover Database for the conterminous United States—Representing a decade of land cover change information, *Photogram. Eng. Rem. Sens.*, *81*, 345–354.
- Hörtensteiner, S. (2006), Chlorophyll degradation during senescence, *Ann. Rev. Plant Biol.*, *57*, 55–77, doi:10.1146/annurev-arplant.57.032905.105212.
- Hawk, F. (2009), Changes in Lake Superior ice cover at Bayfield, Wisconsin, *J. Great Lakes Res.*, *35*, 159–162, doi:10.1016/j.jglr.2008.11.002.
- Huff, A., and A. Thomas (2014), Lake Superior climate change impacts and adaptation Prepared for the Lake Superior Lakewide Action and Management Plan – Superior Work Group. [Available at <http://www.epa.gov/glnpo/lakesuperior/index.html>.]
- Hufkens, K., M. A. Friedl, T. F. Keenan, O. Sonnentag, A. Bailey, J. O’Keefe, and A. D. Richardson (2012), Ecological impacts of a widespread frost event following early spring leaf-out, *Global Change Biol.*, *18*, 2365–2377, doi:10.1111/j.1365-2486.2012.02712.x.
- Hunter, A. F., and M. J. Lechowicz (1992), Predicting the timing of budburst in temperate trees, *J. Appl. Ecol.*, *29*, 597–604, doi:10.2307/2404467.
- Hwang, T., L. E. Band, C. F. Miniati, C. Song, P. V. Bolstad, J. M. Vose, and J. P. Love (2014), Divergent phenological response to hydroclimate variability in forested mountain watersheds, *Global Change Biol.*, *20*, 2580–2595, doi:10.1111/gcb.12556.
- Ireland, L. C., D. Adams, R. Alig, C. J. Betz, C.-C. Chen, M. Hutchins, B. A. McCarl, K. Skog, and B. L. Sohngen (2001), Assessing socioeconomic impacts of climate change on US forests, wood-product markets, and forest recreation, *BioScience*, *51*, 753–764, doi:10.1641/0006-3568(2001)051[0753:asiocc]2.0.co;2.
- Jeong, S.-J., C.-H. Ho, H.-J. Gim, and M. E. Brown (2011), Phenology shifts at start vs. end of growing season in temperate vegetation over the Northern Hemisphere for the period 1982–2008, *Global Change Biol.*, *17*, 2385–2399, doi:10.1111/j.1365-2486.2011.02397.x.
- Jeong, S.-J., D. Medvigy, E. Shevliakova, and S. Malyshev (2013), Predicting changes in temperate forest budburst using continental-scale observations and models, *Geophys. Res. Lett.*, *40*, 359–364, doi:10.1029/2012GL054431.
- Jin, S., L. Yang, P. Danielson, C. Homer, J. Fry, and G. Xian (2013), A comprehensive change detection method for updating the National Land Cover Database to circa 2011, *Remote Sens. Environ.*, *132*, 159–175, doi:10.1016/j.rse.2013.01.012.
- Jo, H.-S., S.-W. Yeh, and B. P. Kirtman (2014), Role of the western tropical Pacific in the North Pacific regime shift in the winter of 1998/1999, *J. Geophys. Res. Oceans*, *119*, 6161–6170, doi:10.1002/2013JC009527.
- Johnson, S. L., and H. G. Stefan (2006), Indicators of climate warming in Minnesota: Lake ice covers and snowmelt runoff, *Clim. Change*, *75*, 421–453, doi:10.1007/s10584-006-0356-0.
- Jolly, W. M., R. Nemani, and S. W. Running (2005), A generalized, bioclimatic index to predict foliar phenology in response to climate, *Global Change Biol.*, *11*, 619–632, doi:10.1111/j.1365-2486.2005.00930.x.
- Keenan, T. F., and A. D. Richardson (2015), The timing of autumn senescence is affected by the timing of spring phenology: Implications for predictive models, *Global Change Biol.*, *21*, 2634–2641, doi:10.1111/gcb.12890.
- Kikuzawa, K., Y. Onoda, I. J. Wright, and P. B. Reich (2013), Mechanisms underlying global temperature-related patterns in leaf longevity, *Global Ecol. Biogeogr.*, *22*, 982–993, doi:10.1111/gcb.12042.
- Killingbeck, K. T. (1996), Nutrients in senesced leaves: Keys to the search for potential resorption and resorption proficiency, *Ecology*, *77*, 1716–1727, doi:10.2307/2265777.
- Kim, Y., J. S. Kimball, K. Didan, and G. M. Henebry (2014), Response of vegetation growth and productivity to spring climate indicators in the conterminous United States derived from satellite remote sensing data fusion, *Agric. For. Meteorol.*, *194*, 132–143, doi:10.1016/j.agrformet.2014.04.001.
- Körner, C. (2015), Paradigm shift in plant growth control, *Curr. Opin. Plant Biol.*, *25*, 107–114, doi:10.1016/j.pbi.2015.05.003.
- Koster, R. D., G. K. Walker, G. J. Collatz, and P. E. Thornton (2014), Hydroclimatic controls on the means and variability of vegetation phenology and carbon uptake, *J. Clim.*, *27*, 5632–5652, doi:10.1175/JCLI-D-13-00477.1.
- Kunkel, K. E., D. R. Easterling, K. Hubbard, and K. Redmond (2004), Temporal variations in frost-free season in the United States: 1895–2000, *Geophys. Res. Lett.*, *31*, L03201, doi:10.1029/2003GL018624.
- Leathers, D. J., B. Yarnal, and M. A. Palecki (1991), The Pacific/North American teleconnection pattern and United States climate. Part I: Regional temperature and precipitation associations, *J. Clim.*, *4*, 517–528, doi:10.1175/1520-0442(1991)004<0517:TPATPA>2.0.CO;2.
- Lechowicz, M. J. (1984), Why do temperate deciduous trees leaf out at different times? Adaptation and ecology of forest communities, *Am. Naturalist*, *124*, 821–842, doi:10.1086/284319.
- Li, X., S. Zhong, X. Bian, W. E. Heilman, Y. Luo, and W. Dong (2010), Hydroclimate and variability in the Great Lakes region as derived from the North American Regional Reanalysis, *J. Geophys. Res.*, *115*, D12104, doi:10.1029/2009JD012756.
- Lu, X., D. W. Kicklighter, J. M. Melillo, J. M. Reilly, and L. Xu (2015), Land carbon sequestration within the conterminous United States: Regional- and state-level analyses, *J. Geophys. Res. Biogeosci.*, *120*, 379–398, doi:10.1002/2014JG002818.
- Luo, D., and J. Cha (2012), The North Atlantic oscillation and the North Atlantic jet variability: Precursors to NAO regimes and transitions, *J. Atmos. Sci.*, *69*, 3763–3787, doi:10.1175/JAS-D-12-098.1.
- Mallard, M. S., C. G. Nolte, O. R. Bullock, T. L. Spero, and J. Gula (2014), Using a coupled lake model with WRF for dynamical downscaling, *J. Geophys. Res. Atmos.*, *119*, 7193–7208, doi:10.1002/2014JD021785.
- Marchin, R., H. Zeng, and W. Hoffmann (2010), Drought-deciduous behavior reduces nutrient losses from temperate deciduous trees under severe drought, *Oecologia*, *163*, 845–854, doi:10.1007/s00442-010-1614-4.
- Marchin, R. M., C. F. Salk, W. A. Hoffmann, and R. R. Dunn (2015), Temperature alone does not explain phenological variation of diverse temperate plants under experimental warming, *Global Change Biol.*, *21*, 3138–3151, doi:10.1111/gcb.12919.
- McCabe, G. J., J. L. Betancourt, and S. Feng (2015), Variability in the start, end, and length of frost-free periods across the conterminous United States during the past century, *Int. J. Climatol.*, doi:10.1002/joc.4315.

- McPhaden, M. J., S. E. Zebiak, and M. H. Glantz (2006), ENSO as an integrating concept in Earth science, *Science*, *314*, 1740–1745, doi:10.1126/science.1132588.
- Menne, M. J., I. Durre, R. S. Vose, B. E. Gleason, and T. G. Houston (2012), An overview of the Global Historical Climatology Network-Daily database, *J. Atmos. Oceanic Technol.*, *29*, 897–910, doi:10.1175/JTECH-D-11-00103.1.
- Millard, P., M. Sommerkorn, and G.-A. Grelet (2007), Environmental change and carbon limitation in trees: A biochemical, ecophysiological and ecosystem appraisal, *New Phytol.*, *175*, 11–28, doi:10.1111/j.1469-8137.2007.02079.x.
- Mishra, V., and K. A. Cherkauer (2011), Influence of cold season climate variability on lakes and wetlands in the Great Lakes region, *J. Geophys. Res.*, *116*, D12111, doi:10.1029/2010JD015063.
- Mishra, V., K. A. Cherkauer, and L. C. Bowling (2010), Parameterization of lakes and wetlands for energy and water balance studies in the Great Lakes region, *J. Hydrometeorol.*, *11*, 1057–1082, doi:10.1175/2010JHM1207.1.
- Mishra, V., K. A. Cherkauer, and L. C. Bowling (2011a), Changing thermal dynamics of lakes in the Great Lakes region: Role of ice cover feedbacks, *Global Planet. Change*, *75*, 155–172, doi:10.1016/j.gloplacha.2010.11.003.
- Mishra, V., K. A. Cherkauer, L. C. Bowling, and M. Huber (2011b), Lake ice phenology of small lakes: Impacts of climate variability in the Great Lakes region, *Global Planet. Change*, *76*, 166–185, doi:10.1016/j.gloplacha.2011.01.004.
- Morin, X., M. J. Lechowicz, C. Augspurger, J. O’Keefe, D. Viner, and I. Chuine (2009), Leaf phenology in 22 North American tree species during the 21st century, *Global Change Biol.*, *15*, 961–975, doi:10.1111/j.1365-2486.2008.01735.x.
- Munné-Bosch, S., and L. Alegre (2004), Die and let live: Leaf senescence contributes to plant survival under drought stress, *Funct. Plant Biol.*, *31*, 203–216, doi:10.1071/FP03236.
- Murray, M. B., M. G. R. Cannell, and R. I. Smith (1989), Date of budburst of fifteen tree species in Britain following climatic warming, *J. Appl. Ecol.*, *26*, 693–700, doi:10.2307/2404093.
- Nemani, R. R., C. D. Keeling, H. Hashimoto, W. M. Jolly, S. C. Piper, C. J. Tucker, R. B. Myneni, and S. W. Running (2003), Climate-driven increases in global terrestrial net primary production from 1982 to 1999, *Science*, *300*, 1560–1563, doi:10.1126/science.1082750.
- Nie, J., P. Wang, W. Yang, and B. Tan (2008), Northern hemisphere storm tracks in strong AO anomaly winters, *Atmos. Sci. Lett.*, *9*, 153–159, doi:10.1002/asl.186.
- Niinemets, Ü., and Ü. Tamm (2005), Species differences in timing of leaf fall and foliage chemistry modify nutrient resorption efficiency in deciduous temperate forest stands, *Tree Physiol.*, *25*, 1001–1014, doi:10.1093/treephys/25.8.1001.
- Ning, L., and R. S. Bradley (2014), Winter precipitation variability and corresponding teleconnections over the northeastern United States, *J. Geophys. Res. Atmos.*, *119*, 7931–7945, doi:10.1002/2014JD021591.
- Noormets, A., S. G. McNulty, J. L. DeForest, G. Sun, Q. Li, and J. Chen (2008), Drought during canopy development has lasting effect on annual carbon balance in a deciduous temperate forest, *New Phytol.*, *179*, 818–828, doi:10.1111/j.1469-8137.2008.02501.x.
- Notaro, M., V. Bennington, and S. Vavrus (2015), Dynamically downscaled projections of lake-effect snow in the Great Lakes basin, *J. Clim.*, *28*, 1661–1684, doi:10.1175/JCLI-D-14-00467.1.
- Nowacki, G. J., and M. D. Abrams (2008), The demise of fire and “mesophication” of forests in the eastern United States, *BioScience*, *58*, 123–138, doi:10.1641/b580207.
- O’Reilly, C. M., et al. (2015), Rapid and highly variable warming of lake surface waters around the globe, *Geophys. Res. Lett.*, *42*, 10,773–10,781, doi:10.1002/2015GL066235.
- Omerik, J. M. (2004), Perspectives on the nature and definitions of ecological regions, *Environ. Manage.*, *34*(Suppl. 1), s27–s38, doi:10.1007/s00267-003-5197-2.
- Omerik, J. M., S. S. Chapman, R. A. Lillie, and R. T. Dumke (2000), Ecoregions of Wisconsin, *Trans. Wisc. Acad. Sci. Arts Lett.*, *88*, 77–103.
- Pagter, M., U. B. Andersen, and L. Andersen (2015), Winter warming delays dormancy release, advances budburst, alters carbohydrate metabolism and reduces yield in a temperate shrub, *AoB Plants*, *7*, plv024, doi:10.1093/aobpla/plv024.
- Parida, B. R., and W. Buermann (2014), Increasing summer drying in North American ecosystems in response to longer nonfrozen periods, *Geophys. Res. Lett.*, *41*, 5476–5483, doi:10.1002/2014GL060495.
- Payer, M., N. F. Laird, R. J. Maliawco Jr., and E. G. Hoffman (2011), Surface fronts, troughs, and baroclinic zones in the Great Lakes region, *Weather Forecast.*, *26*, 555–563, doi:10.1175/WAF-D-10-05018.1.
- Peings, Y., and G. Magnusdottir (2014), Forcing of the wintertime atmospheric circulation by the multidecadal fluctuations of the North Atlantic Ocean, *Environ. Res. Lett.*, *9*, 034018, doi:10.1088/1748-9326/9/3/034018.
- Peñuelas, J., T. Rutishauser, and I. Filella (2009), Phenology feedbacks on climate change, *Science*, *324*, 887–888, doi:10.1126/science.1173004.
- Peters, M. P., L. R. Iverson, and S. N. Matthews (2015), Long-term droughtiness and drought tolerance of eastern US forests over five decades, *For. Ecol. Manage.*, *345*, 56–64, doi:10.1016/j.foreco.2015.02.022.
- Peterson, A. G., and J. T. Abatzoglou (2014), Observed changes in false springs over the contiguous United States, *Geophys. Res. Lett.*, *41*, 2156–2162, doi:10.1002/2014GL059266.
- Peterson, T. C., M. P. Hoerling, P. A. Stott, and S. C. Herring (2013), Explaining extreme events of 2012 from a climate perspective, *Bull. Am. Meteorol. Soc.*, *94*, S1–S74, doi:10.1175/BAMS-D-13-00085.1.
- Peterson, W. T., and F. B. Schwing (2003), A new climate regime in Northeast Pacific ecosystems, *Geophys. Res. Lett.*, *30*(17), 1896, doi:10.1029/2003GL017528.
- Piccolroaz, S., M. Toffolon, and B. Majone (2015), The role of stratification on lakes’ thermal response: The case of Lake Superior, *Water Resour. Res.*, *51*, 7878–7894, doi:10.1002/2014WR016555.
- Plank, C., and B. Shuman (2009), Drought-driven changes in lake areas and their effects on the surface energy balance of Minnesota’s lake-dotted landscape, *J. Clim.*, *22*, 4055–4065, doi:10.1175/2009JCLI1978.1.
- Pope, K. S., V. Dose, D. Da Silva, P. H. Brown, C. A. Leslie, and T. M. DeJong (2013), Detecting nonlinear response of spring phenology to climate change by Bayesian analysis, *Global Change Biol.*, *19*, 1518–1525, doi:10.1111/gcb.12130.
- Prein, A. F., et al. (2015), A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges, *Rev. Geophys.*, *53*, 323–361, doi:10.1002/2014RG000475.
- Reyer, C. P. O., et al. (2013), A plant’s perspective of extremes: Terrestrial plant responses to changing climatic variability, *Global Change Biol.*, *19*, 75–89, doi:10.1111/gcb.12023.
- Richardson, A. D., T. F. Keenan, M. Migliavacca, Y. Ryu, O. Sonnentag, and M. Toomey (2013), Climate change, phenology, and phenological control of vegetation feedbacks to the climate system, *Agric. Forest Meteorol.*, *169*, 156–173, doi:10.1016/j.agrformet.2012.09.012.
- Rigby, J. R., and A. Porporato (2008), Spring frost risk in a changing climate, *Geophys. Res. Lett.*, *35*, L12703, doi:10.1029/2008GL033955.
- Rittenhouse, C. D., and A. R. Rissman (2015), Changes in winter conditions impact forest management in north temperate forests, *J. Environ. Manage.*, *149*, 157–167, doi:10.1016/j.jenvman.2014.10.010.

- Roberts, A. M. I., C. Tansey, R. J. Smithers, and A. B. Phillimore (2015), Predicting a change in the order of spring phenology in temperate forests, *Global Change Biol.*, *21*, 2603–2611, doi:10.1111/gcb.12896.
- Rodionov, S., and R. Assel (2000), Atmospheric teleconnection patterns and severity of winters in the Laurentian Great Lakes basin, *Atmos.-Ocean*, *38*, 601–635, doi:10.1080/07055900.2000.9649661.
- Rodionov, S., and R. Assel (2001), A new look at the Pacific/North American Index, *Geophys. Res. Lett.*, *28*, 1519–1522, doi:10.1029/2000GL012185.
- Rodionov, S., and R. A. Assel (2003), Winter severity in the Great Lakes region: A tale of two oscillations, *Clim. Res.*, *24*, 19–31, doi:10.3354/cr024019.
- Rohde, A., and R. P. Bhalerao (2007), Plant dormancy in the perennial context, *Trends Plant Sci.*, *12*, 217–223, doi:10.1016/j.tplants.2007.03.012.
- Rohli, R. V., A. J. Vega, M. R. Binkley, S. D. Britton, H. E. Heckman, J. M. Jenkins, Y. Ono, and D. E. Sheeler (1999), Surface and 700 hPa atmospheric circulation patterns for the Great Lakes basin and eastern North America and relationship to atmospheric teleconnections, *J. Great Lakes Res.*, *25*, 45–60, doi:10.1016/S0380-1330(99)70716-1.
- Rydzik, M., and A. R. Desai (2014), Relationship between snow extent and midlatitude disturbance centers, *J. Clim.*, *27*, 2971–2982, doi:10.1175/JCLI-D-12-00841.1.
- Saxe, H., M. G. R. Cannell, Ø. Johnsen, M. G. Ryan, and G. Vourlitis (2001), Tree and forest functioning in response to global warming, *New Phytol.*, *149*, 369–399, doi:10.1046/j.1469-8137.2001.00057.x.
- Schaetzl, R. J., B. D. Knapp, and S. A. Isard (2005), Modeling soil temperatures and the mesic-frigid boundary in the central Great Lakes region, 1951–2000, *Soil Sci. Soc. Am. J.*, *69*, 2033–2040, doi:10.2136/sssaj2004.0349.
- Schenker, G., A. Lenz, C. Körner, and G. Hoch (2014), Physiological minimum temperatures for root growth in seven common European broad-leaved tree species, *Tree Physiol.*, *34*, 302–313, doi:10.1093/treephys/tpu003.
- Schwartz, M. D., R. Ahas, and A. Aasa (2006), Onset of spring starting earlier across the Northern Hemisphere, *Global Change Biol.*, *12*, 343–351, doi:10.1111/j.1365-2486.2005.01097.x.
- Schwartz, M. D., J. M. Hanes, and L. Liang (2013), Comparing carbon flux and high-resolution spring phenological measurements in a northern mixed forest, *Ag. Forest Meteorol.*, *169*, 136–147, doi:10.1016/j.agrformet.2012.10.014.
- Scott, R. W., and F. A. Huff (1996), Impacts of the Great Lakes on regional climate conditions, *J. Great Lakes Res.*, *22*, 845–863, doi:10.1016/S0380-1330(96)71006-7.
- Seki, M., T. Yoshida, and T. Takada (2015), A general method for calculating the optimal leaf longevity from the viewpoint of carbon economy, *J. Math. Biol.*, *71*, 669–690, doi:10.1007/s00285-014-0830-7.
- Serbin, S. P., and C. J. Kucharik (2009), Spatiotemporal mapping of temperature and precipitation for the development of a multidecadal climatic dataset for Wisconsin, *J. Appl. Meteorol. Climatol.*, *48*, 742–757, doi:10.1175/2008JAMC1986.1.
- Skaggs, K. E., and S. Irmak (2012), Long-term trends in air temperature distribution and extremes, growing degree-days, and spring and fall frosts for climate impact assessments on agricultural practices in Nebraska, *J. Appl. Meteorol. Climatol.*, *51*, 2060–2073, doi:10.1175/JAMC-D-11-0146.1.
- Small, D., and S. Islam (2009), A synoptic view of trends and decadal variations in autumn precipitation across the United States from 1948 to 2004, *J. Geophys. Res.*, *114*, D10102, doi:10.1029/2008JD011579.
- Small, D., S. Islam, and M. Barlow (2010), The impact of a hemispheric circulation regime on fall precipitation over North America, *J. Hydrometeorol.*, *11*, 1182–1189, doi:10.1175/2010JHM1273.1.
- Sobolowski, S., G. Gong, and M. Ting (2010), Modeled climate state and dynamic responses to anomalous North American snow cover, *J. Clim.*, *23*, 785–799, doi:10.1175/2009JCLI3219.1.
- Sousounis, P. J., and E. K. Grover (2002), Potential future weather patterns over the Great Lakes region, *J. Great Lakes Res.*, *28*, 496–520, doi:10.1016/S0380-1330(02)70602-3.
- Teskey, R. O., B. C. Bongarten, B. M. Cregg, P. M. Dougherty, and T. C. Hennessey (1987), Physiology and genetics of tree growth response to moisture and temperature stress: An examination of the characteristics of loblolly pine (*Pinus taeda* L.), *Tree Physiol.*, *3*, 41–61, doi:10.1093/treephys/3.1.41.
- Thompson, A. J., and S. M. Moncrieff (1982), Prediction of bud burst in Douglas fir by degree-day accumulation, *Can. J. For. Res.*, *12*, 448–452, doi:10.1139/x82-069.
- Trenberth, K. E., G. W. Branstator, and P. A. Arkin (1988), Origins of the 1988 North American drought, *Science*, *242*, 1640–1645, doi:10.1126/science.242.4886.1640.
- Trenberth, K. E., G. W. Branstator, D. Karoly, A. Kumar, N.-C. Lau, and C. Ropelewski (1998), Progress during TOGA in understanding and modeling global teleconnections associated with tropical sea surface temperatures, *J. Geophys. Res.*, *103*, 14,291–14,324, doi:10.1029/97JC01444.
- Trudgill, D. L., A. Honek, D. Li, and N. M. van Straalen (2005), Thermal time—Concepts and utility, *Annals Appl. Biol.*, *146*, 1–14, doi:10.1111/j.1744-7348.2005.04088.x.
- Twine, T. E., and C. J. Kucharik (2009), Climate impacts on net primary productivity trends in natural and managed ecosystems of the central and eastern United States, *Agric. For. Meteorol.*, *149*, 2143–2161, doi:10.1016/j.agrformet.2009.05.012.
- U.S. Environmental Protection Agency (2011), “Level III and IV ecoregions of the continental United States.” U.S. EPA, National Health and Environmental Effects Research Laboratory, Corvallis, Oregon. [Available at http://www.epa.gov/wed/pages/ecoregions/level_iii_iv.htm].
- Van Cleave, K., J. D. Lenters, J. Wang, and E. M. Verhamme (2014), A regime shift in Lake Superior ice cover, evaporation, and water temperature following the warm El Niño winter of 1997–1998, *Limnol. Oceanogr.*, *59*, 1889–1898, doi:10.4319/lo.2014.59.6.1889.
- Viherä-Aarnio, A., S. Soutinen, J. Partanen, and R. Häkkinen (2014), Internal development of vegetative buds of Norway spruce trees in relation to accumulated chilling and forcing temperatures, *Tree Physiol.*, *24*, 547–556, doi:10.1093/treephys/tpu038.
- Vitasse, Y., A. Lenz, and C. Körner (2014), The interaction between freezing tolerance and phenology in temperate deciduous trees, *Front. Plant Sci.*, *5*, 541, doi:10.3389/fpls.2014.00541.
- Voelker, S. L., P.-E. Noiro-Cosson, M. C. Stambaugh, E. R. McMurry, F. C. Meinzer, B. Lachenbruch, and R. P. Guyette (2012), Spring temperature responses of oaks are synchronous with North Atlantic conditions during the last deglaciation, *Ecol. Monogr.*, *82*, 169–187, doi:10.1890/11-0848.1.
- Wang, J., X. Bai, H. Hu, A. Clites, M. Colton, and B. Lofgren (2012), Temporal and spatial variability of Great Lakes ice cover, 1973–2010, *J. Clim.*, *25*, 1318–1329, doi:10.1175/2011JCLI4066.1.
- Watras, C. J., J. S. Read, K. D. Holman, Z. Liu, Y.-Y. Song, A. J. Watras, S. Morgan, and E. H. Stanley (2014), Decadal oscillation of lakes and aquifers in the upper Great Lakes region of North America: Hydroclimatic implications, *Geophys. Res. Lett.*, *41*, 456–462, doi:10.1002/2013GL058679.
- Weaver, S. J., A. Ruiz-Barradas, and S. Nigam (2009), Pentad evolution of the 1988 drought and 1993 flood over the Great Plains: An NARR perspective on the atmospheric and terrestrial water balance, *J. Clim.*, *22*, 5366–5384, doi:10.1175/2009JCLI2684.1.

- Wheeler, G. A., E. J. Cushing, E. Gorham, T. Morley, and G. B. Ownbey (1992), A major floristic boundary in Minnesota: An analysis of 280 taxa occurring in the western and southern portions of the state, *Can. J. Bot.*, *70*, 319–333, doi:10.1139/b92-043.
- White, M. A., P. E. Thornton, and S. W. Running (1997), A continental phenology model for monitoring vegetation responses to interannual climatic variability, *Global Biogeochem. Cycles*, *11*, 217–234, doi:10.1029/97GB00330.
- Williams, C. M., H. A. L. Henry, and B. J. Sinclair (2014), Cold truths: How winter drives responses of terrestrial organisms to climate change, *Biol. Rev.*, *90*, 214–235, doi:10.1111/brv.12105.
- Wise, E. K., M. L. Wrzesien, M. P. Dannenberg, and D. L. McGinnis (2015), Cool-season precipitation patterns associated with teleconnection interactions in the United States, *J. Appl. Meteorol. Climatol.*, *54*, 494–505, doi:10.1175/JAMC-D-14-0040.1.
- Wright, D. M., D. J. Posselt, and A. L. Steiner (2013), Sensitivity of lake-effect snowfall to lake ice cover and temperature in the Great Lakes region, *Mon. Weather Rev.*, *141*, 670–689, doi:10.1175/MWR-D-12-00038.1.
- Xie, Y., K. F. Ahmed, J. M. Allen, A. M. Wilson, and J. A. Silander Jr. (2015), Green-up of deciduous forest communities of northeastern North America in response to climate variation and climate change, *Lands. Ecol.*, *30*, 109–123, doi:10.1007/s10980-014-0099-7.
- Yang, L. H., K. F. Edwards, J. E. Byrnes, J. L. Bastow, A. N. Wright, and K. O. Spence (2010), A meta-analysis of resource pulse–consumer interactions, *Ecol. Monogr.*, *80*, 125–151, doi:10.1890/08-1996.1.
- Yebra, M., P. E. Dennison, E. Chuvieco, D. Riaño, P. Zylstra, E. R. Hunt Jr., F. M. Danson, Y. Qi, and S. Jurdao (2013), A global review of remote sensing of live fuel moisture content for fire danger assessment: Moving towards operational products, *Remote Sens. Environ.*, *136*, 455–468, doi:10.1016/j.rse.2013.05.029.
- Yu, L., S. Zhong, X. Bian, W. E. Heilman, and J. A. Andresen (2014), Temporal and spatial variability of frost-free seasons in the Great Lakes region of the United States, *Int. J. Climatol.*, *34*, 3499–3514, doi:10.1002/joc.3923.